

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer |
|------------------------|--|
| Catalog Number: | B6117S |
| Concentration: | 10 X Concentrate |
| Packaging Lot Number: | 10098905 |
| Expiration Date: | 12/2022 |
| Storage Temperature: | -20°C |
| Specification Version: | PS-B6117S v2.0 |
| Composition (1X): | 20 mM Tris-HCl, 12 mM (NH4)2SO4, 5 mM MgCl2, 0.16 mM β-NAD, (pH 7.5 @ 25°C) |

| NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer Component List | | | | |
|---|---|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| B6117SVIAL | NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer | 10095144 | Pass | |

| Assay Name/Specification | Lot # 10098905 |
|--|----------------|
| Endonuclease Activity (Nicking, Buffer) A 50 μ l reaction in 1X NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer containing 1 μ g of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Phosphatase Activity (pNPP, Buffer) A 200 μl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 μl NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis. | Pass |
| RNase Activity (Buffer) A 10 µl reaction in 1X NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

mostin_ /

Christine Sumner Production Scientist 29 Jan 2021

Michae on. 1

Michael Tonello Packaging Quality Control Inspector 29 Jan 2021

