

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer
Catalog Number:	B6117S
Concentration:	10 X Concentrate
Packaging Lot Number:	10139420
Expiration Date:	02/2023
Storage Temperature:	-20°C
Specification Version:	PS-B6117S v2.0
Composition (1X):	20 mM Tris-HCl, 12 mM (NH4)2SO4, 5 mM MgCl2, 0.16 mM β-NAD, (pH 7.5 @ 25°C)

NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B6117SVIAL	NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer	10104490	Pass	

Assay Name/Specification	Lot # 10139420
RNase Activity (Buffer) A 10 µl reaction in 1X NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 μ I reaction in 1X NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer containing 1 μ g of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP, Buffer) A 200 μl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 μl NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X NEBNext® Second Strand Synthesis (dNTP-free) Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

mostin_ /

Christine Sumner Production Scientist 31 Jan 2022

Michae on. 1

Michael Tonello Packaging Quality Control Inspector 31 Jan 2022

