

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	dam-/dcm- Competent E. coli
Catalog Number:	C2925I
Packaging Lot Number:	10196214
Expiration Date:	05/2024
Storage Temperature:	-80°C
Specification Version:	PS-C2925H/I v1.0

dam-/dcm- Competent E. coli Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N3041AVIAL	pUC19 Vector	10181834	Pass	
C2925IVIAL	dam ⁻ /dcm ⁻ Competent E. coli	10176318	Pass	
B9020SVIAL	SOC Outgrowth Medium	10174270	Pass	

5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass antibiotic Resistance (Nitrofurantoin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 9 µl of untransformed dam-/dcm- Competent E	Assay Name/Specification	Lot # 10196214
ontaining Chloramphenicol will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Resistance (Nitrofurantoin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Resistance (Streptomycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Streptomycin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Ampicillin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.Pass	Antibiotic Resistance (Chloramphenicol)	Pass
Antibiotic Resistance (Nitrofurantoin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Resistance (Streptomycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Streptomycin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Ampicillin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePass		
 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Nitrofurantoin will form colonies after incubation for 16 hours at 37°C. Antibiotic Resistance (Streptomycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Streptomycin will form colonies after incubation for 16 hours at 37°C. Antibiotic Sensitivity (Ampicillin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C. Antibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C. Antibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate 		
ontaining Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Resistance (Streptomycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Streptomycin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Ampicillin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePassAntibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePass	Antibiotic Resistance (Nitrofurantoin)	Pass
Antibiotic Resistance (Streptomycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Streptomycin will form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Ampicillin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePassAntibiotic Sensitivity (Kanamycin) 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePass		
 5 μl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Streptomycin will form colonies after incubation for 16 hours at 37°C. Antibiotic Sensitivity (Ampicillin) 5 μl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C. Antibiotic Sensitivity (Kanamycin) 5 μl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass 		
Antibiotic Sensitivity (Ampicillin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePassAntibiotic Sensitivity (Kanamycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePass	Antibiotic Resistance (Streptomycin)	Pass
Antibiotic Sensitivity (Ampicillin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C.PassAntibiotic Sensitivity (Kanamycin)Pass5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth platePass		
5 μl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C. Intibiotic Sensitivity (Kanamycin) 5 μl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate		
ontaining Ampicillin will not form colonies after incubation for 16 hours at 37°C. Pass Antibiotic Sensitivity (Kanamycin) Pass 5 μl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass	Antibiotic Sensitivity (Ampicillin)	Pass
Antibiotic Sensitivity (Kanamycin) Pass 5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate Pass		
5 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate		
	Antibiotic Sensitivity (Kanamycin)	Pass
	Antibiotic Sensitivity (Spectinomycin)	Pass
	15 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at	





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10196214
37°C.	
Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	Pass
Phage Resistance (ϕ 80) 15 µl of untransformed dam-/dcm- Competent E. coli streaked onto a Rich Broth plate does not support plaque formation by phage ϕ 80 after incubation for 16 hours at 37°C.	Pass
Transformation Efficiency 50 μ l of dam-/dcm- Competent E. coli cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e6 cfu/ μ g of DNA.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Q Pr duction Scientist 14 Feb 2023

M

Nick Privitera Packaging Quality Control Inspector 07 Jun 2023

