

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

NEB® Turbo Electrocompetent E. coli
C2986K
0241803
03/2018
03/2019
-80°C
PS-C2986K v1.0
07 Feb 2018

Assay Name/Specification (minimum release criteria)	Lot #0241803
Antibiotic Resistance (Nitrofurantoin) - 15 μ l of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Ampicillin) - 15 μ l of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Chloramphenicol) - 15 μ l of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Kanamycin) - 15 μ l of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Spectinomycin) - 15 µl of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Streptomycin) - 15 μ l of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37° C.	Pass
Antibiotic Sensitivity (Tetracycline) - 15 μ l of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a LB or Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37° C.	Pass
Blue-White Screening (α-complementation, Competent Cells) - NEB® Turbo Electrocompetent <i>E. coli</i> were shown to be suitable for blue/white screening by α -complementation of the β -galactosidase gene using pUC19.	Pass



C2986K Lot: 0241803 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0241803
Phage Resistance (Φ 80) - 15 µl of untransformed NEB® Turbo Electrocompetent <i>E. coli</i> streaked onto a Rich Broth plate does not support plaque formation by phage Φ 80 after incubation for 16 hours at 37°C.	Pass
Transformation Efficiency - 25 μ l of NEB® Turbo Electrocompetent <i>E. coli</i> cells were transformed with 10 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37° C resulted in >1 x 10e10 cfu/µg of DNA.	Pass

un otunon

Authorized by Derek Robinson 07 Feb 2018



Inspected by Quiting Ren 10 Apr 2018