

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: NEB® 5-alpha Competent E. coli (Subcloning Efficiency)

Catalog Number: C2988J
Packaging Lot Number: 10163364
Expiration Date: 08/2023
Storage Temperature: -80°C

Specification Version: PS-C2988J v1.0

NEB® 5-alpha Competent E. coli (Subcloning Efficiency) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
C2988JVIAL	NEB® 5-alpha Competent E. coli (Subcloning Efficiency)	10145023	Pass	

Assay Name/Specification	Lot # 10163364
Transformation Efficiency 50 μl of NEB® 5-alpha Competent E. coli (Subcloning Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e6 cfu/μg of DNA.	Pass
Phage Resistance (φ 80) 15 μl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.	Pass
Blue-White Screening (α -complementation, Competent Cells) NEB® 5-alpha Competent E. coli (Subcloning Efficiency) were shown to be suitable for blue/white screening by α -complementation of the β -galactosidase gene using pUC19.	Pass
Antibiotic Sensitivity (Nitrofurantoin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Chloramphenicol) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.	Pass



C2988J / Lot: 10163364

Page 1 of 2

Assay Name/Specification	Lot # 10163364
Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Streptomycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Spectinomycin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed NEB® 5-alpha Competent E. coli (Subcloning Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lixin An

Production Scientist

02 Sep 2022

Corey Rabeau

Packaging Quality Control Inspector

02 Sep 2022



C2988J / Lot: 10163364

Page 2 of 2