

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: LongAmp® Taq 2X Master Mix

Catalog Number: M0287L

Concentration: 2 X Concentrate

Packaging Lot Number: 10084988
Expiration Date: 09/2021
Storage Temperature: -20°C

Specification Version: PS-M0287S/L v2.0

Composition (1X): 60 mM Tris-SO4 (pH 9.1 @ 25°C), 20 mM (NH4)2SO4, 2 mM MgSO4, 0.3 mM

dATP, 0.3 mM dCTP, 0.3 mM dGTP, 0.3 mM dTTP, 3 % Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 125 units/ml LongAmp® Taq DNA

Polymerase

| LongAmp® Taq 2X Master Mix Component List | | | |
|---|----------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M0287SVIAL | LongAmp® Taq 2X Master Mix | 10069425 | Pass |

| Assay Name/Specification | Lot # 10084988 |
|--|----------------|
| PCR Amplification (30 kb Lambda DNA, Master Mix) | Pass |
| A 25 μl reaction in 1X LongAmp® Taq Master Mix and 0.4 μM primers containing 1 ng | |
| Lambda DNA for 28 cycles of PCR amplification results in the expected 30 kb product. | |
| qPCR DNA Contamination (E. coli Genomic) | Pass |
| A minimum of 2.5 units of LongAmp® Taq DNA Polymerase is screened for the presence | |
| of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli | |
| 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA | |
| contamination is ≤ 1 E. coli genome. | |
| | |
| RNase Activity (Extended Digestion) | Pass |
| A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA | |
| and a minimum of 1 µl of LongAmp® Taq 2X Master Mix is incubated at 37°C. After | |
| incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | |
| gor orotrophorotic deling nuorescent detection. | |
| Non-Specific DNase Activity (16 hour, Buffer) | Pass |
| A 50 μl reaction in 1X LongAmp® Taq Master Mix containing 1 μg of T3 or T7 DNA in | |
| addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C | |
| results in a DNA pattern free of detectable nuclease degradation as determined by | |



M0287L / Lot: 10084988 Page 1 of 2 This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Christie Vazquez **Production Scientist**

13 Oct 2020

Josh Hersey

Packaging Quality Control Inspector

13 Oct 2020

