

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Shrimp Alkaline Phosphatase (rSAP)

Catalog Number: M0371L
Concentration: 1,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 µmol

of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml

in 1 minute at 37°C

Packaging Lot Number: 10153911
Expiration Date: 07/2024
Storage Temperature: -20°C

Storage Conditions: 25 mM Tris-HCl , 1 mM MgCl2 , 50 % Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0371S/L v1.0

Shrimp Alkaline Phosphatase (rSAP) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0371LVIAL	Shrimp Alkaline Phosphatase (rSAP)	10153910	Pass
B6004SVIAL	rCutSmart™ Buffer	10154051	Pass

Assay Name/Specification	Lot # 10153911
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 10 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Shrimp Alkaline Phosphatase (rSAP) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Protein Purity Assay (SDS-PAGE)	Pass



M0371L / Lot: 10153911

Page 1 of 2

Assay Name/Specification	Lot # 10153911
Shrimp Alkaline Phosphatase (rSAP) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum of 10 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist

26 Jul 2022

Erin Varney

Packaging Quality Control Inspector

26 Jul 2022



M0371L / Lot: 10153911

Page 2 of 2