

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Shrimp Alkaline Phosphatase (rSAP)

Catalog Number: M0371S
Concentration: 1,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 µmol

of p-Nitrophenyl Phosphate, PNPP in a total reaction volume of 1 ml

in 1 minute at 37°C

Packaging Lot Number: 10111319
Expiration Date: 05/2023
Storage Temperature: -20°C

Storage Conditions: 25 mM Tris-HCl , 1 mM MgCl2 , 50 % Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0371S/L v1.0

Shrimp Alkaline Phosphatase (rSAP) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0371SVIAL	Shrimp Alkaline Phosphatase (rSAP)	10106620	Pass	
B6004SVIAL	rCutSmart™ Buffer	10111602	Pass	

Assay Name/Specification	Lot # 10111319
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 5 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Shrimp Alkaline Phosphatase (rSAP) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Shrimp Alkaline Phosphatase (rSAP) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 10 units of Shrimp Alkaline	Pass



M0371S / Lot: 10111319

Page 1 of 2

Assay Name/Specification	Lot # 10111319
Phosphatase (rSAP) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum of 10 units of Shrimp Alkaline Phosphatase (rSAP) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 15 Jul 2021 Michael Tonello

Packaging Quality Control Inspector

15 Jul 2021



M0371S / Lot: 10111319

Page 2 of 2