

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:RtcB LigaseCatalog Number:M0458SConcentration:15 μMPackaging Lot Number:10068444Expiration Date:02/2022Storage Temperature:-20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

(pH 7.4 @ 25°C)

Specification Version: PS-M0458S v1.0

RtcB Ligase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
N2080AVIAL	GTP	10067100	Pass	
M0458SVIAL	RtcB Ligase	10068443	Pass	
B1761SVIAL	10mM MnCl2	10048443	Pass	
B0790AVIAL	10X RtcB Reaction Buffer	10068468	Pass	

Assay Name/Specification	Lot # 10068444
Endonuclease Activity (Nicking) A 50 µl reaction in RtcB Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 15 pmol of RtcB Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in RtcB Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 15 pmol of RtcB Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (RtcB Ligase) A 20 μl reaction in 1X RtcB Reaction Buffer supplemented with 0.1 mM GTP and 1 mM MnCl2 containing 0.5 μM of a 17 mer 5' FAM-labeled 3' phosphorylated RNA, 0.5 μM of a 30 mer 5' OH RNA and 0.75 μM RtcB Ligase incubated for 1 hour at 37°C results in ≥ 80% ligation as determined by capillary electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) RtcB Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



M0458S / Lot: 10068444

Page 1 of 2

Assay Name/Specification	Lot # 10068444
DNose Activity (Extended Digestion)	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass
and a minimum of 15 pmol of RtcB Ligase is incubated at 37°C. After incubation for	
16 hours, >90% of the substrate RNA remains intact as determined by gel	
electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

Bhairavi Jani Production Scientist

07 May 2020

Jay Minichiello

Packaging Quality Control Inspector

07 May 2020



M0458S / Lot: 10068444 Page 2 of 2