

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: OneTag® Hot Start 2X Master Mix with Standard Buffer

Catalog Number: M0484S

Concentration: 2 X Concentrate

Lot Number: 10042651
Expiration Date: 01/2021
Storage Temperature: -20°C

Specification Version: PS-M0484S/L v1.0

Composition (1X): 20 mM Tris-HCl (pH 8.9 @ 25°C), 22 mM NH4Cl, 22 mM KCl, 1.8 mM

MgCl2, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 25 units/ml

OneTaq® Hot Start DNA Polymerase

OneTaq® Hot Start 2X Master Mix with Standard Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0484SVIAL	OneTaq® Hot Start 2X Master Mix with Standard Buffer	10032983	Pass	

Assay Name/Specification	Lot # 10042651
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq® Hot Start 2X Master Mix with Standard Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
PCR Amplification (Hot Start 2 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.	Pass
PCR Amplification (5 kb Lambda, Master Mix) A 25 μl reaction in 1X OneTaq® Hot Start Master Mix with Standard Buffer and 0.2 μM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X OneTaq® Hot Start Master Mix with Standard Buffer containing	Pass



M0484S / Lot: 10042651

Page 1 of 2

Assay Name/Specification	Lot # 10042651
1 μg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs including [³H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction.	Pass

This product has been tested and shown to be in compliance with all specifications.

Christie Vazquez **Production Scientist**

24 Jan 2019

Josh Hersey

Packaging Quality Control Inspector

13 May 2019



M0484S / Lot: 10042651

Page 2 of 2