

OVERY 240 County Road
JINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pyrophosphatase, inorganic (yeast)

Catalog Number: M2403S
Concentration: 100 U/ml

Unit Definition: One unit is the amount of enzyme that will generate 1 µmol of

phosphate per minute from inorganic pyrophosphate under standard

reaction conditions.

Packaging Lot Number: 10098423
Expiration Date: 02/2023
Storage Temperature: -20°C

Storage Conditions: 100 mM KCI, 20 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

(pH 8.0 @ 25°C)

Specification Version: PS-M2403S/L v2.0

Pyrophosphatase, inorganic (yeast) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M2403SVIAL	Pyrophosphatase, inorganic (yeast)	10098422	Pass	

Assay Name/Specification	Lot # 10098423
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 0.1 unit of Pyrophosphatase, Inorganic (yeast) is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
dNTPase Activity A 0.5 ml reaction in ThermoPol® Reaction Buffer in the presence of 200 μM each dNTPs and a minimum of 1 unit Pyrophosphatase, Inorganic (yeast) incubated for 1 hour at 37°C results in <0.05 μmol of inorganic phosphate from dNTPs as determined by the AAM assay.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 4 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 unit of Pyrophosphatase, Inorganic (yeast) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour)	Pass



M2403S / Lot: 10098423

Page 1 of 2

Assay Name/Specification	Lot # 10098423
A 50 µl reaction in NEBuffer 4 containing 1 µg of Lambda DNA and a minimum of 1 unit of Pyrophosphatase, Inorganic (yeast) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 1 unit of Pyrophosphatase, Inorganic (yeast) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Phosphatase Activity (pNPP) A 100 µl reaction in NEBuffer 3 containing 10 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 1 unit Pyrophosphatase, Inorganic (yeast) incubated for 1 hour at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Timothy Meixsell Production Scientist 05 Mar 2021 Michael Tonello

Packaging Quality Control Inspector

05 Mar 2021

M2403S / Lot: 10098423