

## New England Biolabs Certificate of Analysis

**Product Name:** *Thermolabile USER<sup>®</sup> II Enzyme*

**Catalog Number:** *M5508S*

**Concentration:** *1,000 U/ml*

**Unit Definition:** *One unit is defined as the amount of enzyme required to nick 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base in 15 minutes at 37°C in a total reaction volume of 10 µL in 1X T4 DNA Ligase Buffer.*

**Packaging Lot Number:** *10092492*

**Expiration Date:** *10/2022*

**Storage Temperature:** *-20°C*

**Storage Conditions:** *25 mM KCl, 35 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 15 mM Tris-HCl, 100 µg/ml BSA, 50 % Glycerol, (pH 7.5 @ 25°C)*

**Specification Version:** *PS-M5508S/L v1.0*


Thermolabile USER <sup>®</sup> II Enzyme Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M5508SVIAL	Thermolabile USER <sup>®</sup> II Enzyme	10086076	Pass
B7204SVIAL	CutSmart <sup>®</sup> Buffer	10091029	Pass

Assay Name/Specification	Lot # 10092492
<p><b>Functional Testing (Thermolability, Endonuclease III)</b></p> <p>A 10 µl reaction in CutSmart<sup>®</sup> Buffer containing 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base and 1 unit of Thermolabile USER<sup>®</sup> II Enzyme was incubated for 15 minutes at 37°C followed by heat inactivation for 10 minutes at 65°C. The addition of 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single AP site and incubation for 15 minutes at 37°C followed by 10 minutes at 75°C, results in no cleavage of additional substrate.</p>	Pass
<p><b>Functional Testing (Thermolability, UDG)</b></p> <p>A 10 µl reaction in CutSmart<sup>®</sup> Buffer containing 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base and 1 unit of Thermolabile USER<sup>®</sup> II Enzyme was incubated for 15 minutes at 37°C followed by heat inactivation for 10 minutes at 65°C. The addition of 10 pmol of a 34 mer fluorescently labeled oligonucleotide duplex containing a single uracil base with 20 units of Endonuclease III and incubation for 15 minutes at 37°C followed by 10 minutes at 75°C, results in no cleavage of additional substrate.</p>	Pass

Assay Name/Specification	Lot # 10092492
<p><b>Functional Testing (USER, Transformation assay)</b> A 10 µl reaction in ThermoPol<sup>®</sup> Reaction Buffer containing 20 ng linearized pNEB206A, 100 ng of a 950 bp control PCR product and 1 unit of Thermolabile USER<sup>®</sup> II Enzyme was incubated for 15 minutes at 37°C followed by 15 minutes at 25°C. After transformation into ER2267 chemically-competent cells &gt;95% of colonies contained recombinant plasmid.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 1 unit of Thermolabile USER<sup>®</sup> II Enzyme is screened for the presence of E. coli genomic DNA using SYBR<sup>®</sup> Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Thermolabile USER<sup>®</sup> II Enzyme is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Lauren Higgins  
Production Scientist  
10 Dec 2020



Michael Tonello  
Packaging Quality Control Inspector  
10 Dec 2020