

New England Biolabs Certificate of Analysis


Product Name: dATP Solution
Catalog Number: N0440S
Concentration: 100 mM
Unit Definition: N/A
Lot Number: 10048409
Expiration Date: 06/2021
Storage Temperature: -20°C
Storage Conditions: Supplied in Ultrapure water as a sodium salt (pH 7.5)
Specification Version: PS-N0440S v1.0

| dATP Solution Component List | | | |
|------------------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| N0440SVIAL | dATP Solution | 10044848 | Pass |

| Assay Name/Specification | Lot # 10048409 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 1 µl of dATP Solution incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 µl of dATP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| PCR Amplification (0.5 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dGTP, dCTP, and dTTP and 0.5 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product. | Pass |
| PCR Amplification (2.0 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dGTP, dCTP, and dTTP and 0.5 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product. | Pass |

| Assay Name/Specification | Lot # 10048409 |
|---|----------------|
| <p>PCR Amplification (5.0 kb Lambda, dNTPs) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dATP, dGTP, dCTP, and dTTP and 0.5 µM primers containing 1 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.</p> | Pass |
| <p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 4 µl dATP Solution incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p> | Pass |
| <p>Physical Purity (HPLC) dATP Solution is ≥ 99% pure as determined by HPLC analysis.</p> | Pass |
| <p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of dATP Solution is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.



David Guo
Production Scientist
03 Jul 2019



Michael Tonello
Packaging Quality Control Inspector
15 Jul 2019