

New England Biolabs Certificate of Analysis

Product Name: PNGase F (Glycerol-free), Recombinant
Catalog #: P0709S/L
Concentration: 500,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to remove > 95% of the carbohydrate from 10 µg of denatured RNase B in 1 hour at 37°C in a total reaction volume of 10 µl. (65 NEB units = 1 IUB milliunit).
Lot #: 0021703
Assay Date: 03/2017
Expiration Date: 03/2019
Storage Temp: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0709S/L v1.0
Effective Date: 16 Feb 2016

Assay Name/Specification (minimum release criteria)	Lot #0021703
Glycosidase Activity (Endo F1, F2, H) - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 5,000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (Endo F2, F3) - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Mannosidase) - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β-Xylosidase) - A 10 µl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0021703
<p>Glycosidase Activity (β1-3 Galactosidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β1-4 Galactosidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Glucosidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α-Neuraminidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-2 Fucosidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-2Galβ1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p>Glycosidase Activity (α1-3 Fucosidase) - A 10 μl reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 5,000 of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0021703
Glycosidase Activity (α1-3 Galactosidase) - A 10 μ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-3Gal β 1-4GlcNAc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-3 Mannosidase) - A 10 μ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Galactosidase) - A 10 μ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α1-6 Mannosidase) - A 10 μ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-6Man α 1-6(Man α 1-3)Man-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α-N-Acetylgalactosaminidase) - A 10 μ l reaction in Glyco Buffer 2 containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC) and 5000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Protease Activity (SDS-PAGE) - A 20 μ l reaction in 1X Glyco Buffer 2 containing 24 μ g of a standard mixture of proteins and a minimum of 10,000 units of PNGase F (Glycerol-free), Recombinant incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) - PNGase F (Glycerol-free), Recombinant is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



Authorized by
Derek Robinson
16 Feb 2016



Inspected by
Alicia Bielik
24 Mar 2017

