Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EGCase I P0773S Catalog #: Concentration: 6 units/ml

Unit Definition: One unit of R. triatomea EGCase I is defined as the amount of enzyme required to hydrolyze 1 µmol of ganglioside GM1a per

minute at 37°C.

Lot #: 0011802 Assay Date: 02/2018 Expiration Date: 02/2020 Storage Temp: *-20°C*

Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)

Specification Version: PS-P0773S v1.0 Effective Date: 03 Aug 2017

Assay Name/Specification (minimum release criteria)	Lot #0011802
Glycosidase Activity (Endo F1, F2, H) - A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (Endo F2, F3) - A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (PNGase F) - A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β -Mannosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled β -Mannosidase substrate (Man β 1-4Man β 1-4Man-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β -Xylosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled β -Xylosidase substrate (Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β1-3 Galactosidase) - A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass









New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0011802
Glycosidase Activity ($β1-4$ Galactosidase) - A 10 $μ$ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled $β$ -Galactosidase substrate (Gal $β1-4$ GlcNAc $β1-3$ Gal $β1-4$ Glc -AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β -N-Acetylgalactosaminidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled β -N-Acetylgalactosaminidase substrate (GalNAc β 1-4Gal β 1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (β -N-Acetylglucosaminidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α -Glucosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α -Neuraminidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-2 Fucosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently -labeled α -Fucosidase substrate (Fuc α 1-2Gal β 1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α 1-3 Fucosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently -labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity ($\alpha 1$ -3 Galactosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal $\alpha 1$ -3Gal $\beta 1$ -4GlcNAc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity ($\alpha 1$ -3 Mannosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man $\alpha 1$ -3Man $\beta 1$ -4GlcNAc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass









New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0011802
Glycosidase Activity (α 1-6 Galactosidase) - A 10 μl reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity ($\alpha 1$ -6 Mannosidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man $\alpha 1$ -6(Man $\alpha 1$ -6(Man $\alpha 1$ -3)Man-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Glycosidase Activity (α -N-Acetylgalactosaminidase) - A 10 μ l reaction in EGCase I Buffer containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC) and 6 mU of EGCase I incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.	Pass
Protease Activity (SDS-PAGE) - A 20 μl reaction in 1X EGCase I Buffer containing 24 μg of a standard mixture of proteins and a minimum of 30 mU of EGCase I incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.	Pass
Protein Purity Assay (SDS-PAGE) - EGCase I is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

Authorized by

Derek Robinson
03 Aug 2017







×

Inspected by Saulius Vainauskas 27 Feb 2018