

## New England Biolabs Certificate of Analysis

**Product Name:** XhoI  
**Catalog Number:** R0146L  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) fragments in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10122537  
**Expiration Date:** 05/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R0146S/L v2.0

XhoI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0146LVIAL	XhoI	10107342	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10108731	Pass
B6004SVIAL	rCutSmart™ Buffer	10120518	Pass

Assay Name/Specification	Lot # 10122537
<b>Blue-White Screening (Terminal Integrity)</b> A sample of Litmus 28i vector linearized with a 10-fold excess of XhoI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> XhoI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of pXba DNA with XhoI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with XhoI.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII DNA and a minimum of 100 Units of XhoI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass

