

## New England Biolabs Certificate of Analysis

**Product Name:** ThermoPol<sup>®</sup> Reaction Buffer Pack  
**Catalog Number:** B9004S  
**Concentration:** 10 X Concentrate  
**Packaging Lot Number:** 10122155  
**Expiration Date:** 02/2026  
**Storage Temperature:** -20°C  
**Specification Version:** PS-B9004S v2.0  
**Composition (1X):** 20 mM Tris-HCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 10 mM KCl, 2 mM MgSO<sub>4</sub>, 0.1 % Triton<sup>®</sup>X-100, (pH 8.8 @ 25°C)

ThermoPol <sup>®</sup> Reaction Buffer Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B9004SVIAL	ThermoPol <sup>®</sup> Reaction Buffer Pack	10125244	Pass
B1003SVIAL	Magnesium Sulfate (MgSO <sub>4</sub> ) Solution	10118450	Pass

Assay Name/Specification	Lot # 10122155
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 2X ThermoPol <sup>®</sup> Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 2X ThermoPol <sup>®</sup> Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>PCR Amplification (5 kb Lambda DNA, Buffer)</b> A 50 µl reaction in ThermoPol <sup>®</sup> Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
<b>Phosphatase Activity (pNPP, Buffer)</b> A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl ThermoPol <sup>®</sup> Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass


Assay Name/Specification	Lot # 10122155
<p><b>qPCR DNA Contamination (E. coli Genomic, Buffer)</b> A minimum of 1 µl of ThermoPol® Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of ThermoPol® Reaction Buffer is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>
<p><b>pH (buffers/solutions)</b> The pH of 10X ThermoPol® Reaction Buffer is between pH 8.7 and 8.9 at 25°C.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Christie Vazquez  
Production Scientist  
17 Nov 2021



Michael Tonello  
Packaging Quality Control Inspector  
17 Nov 2021