

New England Biolabs Certificate of Analysis

Product Name: *Apal*
Catalog Number: *R0114S*
Concentration: *50,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10137200*
Expiration Date: *01/2024*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)*
Specification Version: *PS-R0114S/L v2.0*

Apal Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0114SVIAL	Apal	10137117	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10130600	Pass
B6004SVIAL	rCutSmart™ Buffer	10131975	Pass

Assay Name/Specification	Lot # 10137200
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 100 units of Apal incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pXba DNA and 1 µl of Apal incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of Apal incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 50 units of Apal is screened for the presence of E. coli genomic DNA	Pass

Assay Name/Specification	Lot # 10137200
<p>using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	
<p>Protein Purity Assay (SDS-PAGE) Apal is $\geq 95\%$ pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [3H] E. coli DNA and a minimum of 100 units of Apal incubated for 4 hours at 37°C releases $<0.1\%$ of the total radioactivity.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pXba DNA with Apal, $>95\%$ of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, $>95\%$ can be recut with Apal.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
04 Mar 2022



Josh Hersey
Packaging Quality Control Inspector
04 Mar 2022