

## New England Biolabs Certificate of Analysis

**Product Name:** Sall-HF<sup>®</sup>  
**Catalog Number:** R3138L  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10030758  
**Expiration Date:** 12/2020  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 300 µg/ml BSA  
**Specification Version:** PS-R3138S/L v1.0

Sall-HF <sup>®</sup> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3138LVIAL	Sall-HF <sup>®</sup>	10030759	Pass
B7204SVIAL	CutSmart <sup>®</sup> Buffer	10021123	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10021129	Pass


Assay Name/Specification	Lot # 10030758
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of Sall-HF <sup>™</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 50-fold over-digestion of Adenovirus-2 DNA with Sall-HF <sup>™</sup> , >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with Sall-HF <sup>™</sup> .	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of pBR322 DNA and a minimum of 200 Units of Sall-HF <sup>™</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Blue-White Screening (Terminal Integrity)</b> A sample of pUC19 vector linearized with a 10-fold excess of Sall-HF <sup>™</sup> , religated	Pass

Assay Name/Specification	Lot # 10030758
<p>and transformed into an E. coli strain expressing the LacZ beta fragment gene results in &lt;1% white colonies.</p> <p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of Sall-HF™ incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



Tony Spear-Alfonso  
Production Scientist  
08 Nov 2018



Michael Tonello  
Packaging Quality Control Inspector  
06 Dec 2018