

## Taq DNA Ligase Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0208S 003160919091

# B0208S

6.0 ml Taq DNA Ligase Reaction Buffer (10X)

Store at  $-20^{\circ}\text{C}$  Lot: 0031609 Exp: 9/19

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### 1X Taq DNA Ligase Reaction Buffer:

20 mM Tris-HCl  
25 mM potassium acetate  
10 mM magnesium acetate  
10 mM DTT  
1 mM NAD  
0.1% Triton X-100  
pH 7.6 @  $25^{\circ}\text{C}$

## Taq DNA Ligase Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0208S 003160919091

# B0208S

6.0 ml Taq DNA Ligase Reaction Buffer (10X)

Store at  $-20^{\circ}\text{C}$  Lot: 0031609 Exp: 9/19

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### 1X Taq DNA Ligase Reaction Buffer:

20 mM Tris-HCl  
25 mM potassium acetate  
10 mM magnesium acetate  
10 mM DTT  
1 mM NAD  
0.1% Triton X-100  
pH 7.6 @  $25^{\circ}\text{C}$

### Quality Control Assays

**16-Hour Incubation:** A 50  $\mu\text{l}$  reaction containing this reaction buffer at a 1X concentration and 1  $\mu\text{g}$  of HaeIII digested  $\phi\text{X174}$  RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1  $\mu\text{g}$   $\phi\text{X174}$  RF I DNA for 4 hours at  $37^{\circ}\text{C}$  in 50  $\mu\text{l}$  reactions resulted in less than 5% conversion to RF II.



NEW ENGLAND BIOLABS<sup>®</sup> is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

### Quality Control Assays

**16-Hour Incubation:** A 50  $\mu\text{l}$  reaction containing this reaction buffer at a 1X concentration and 1  $\mu\text{g}$  of HaeIII digested  $\phi\text{X174}$  RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1  $\mu\text{g}$   $\phi\text{X174}$  RF I DNA for 4 hours at  $37^{\circ}\text{C}$  in 50  $\mu\text{l}$  reactions resulted in less than 5% conversion to RF II.



NEW ENGLAND BIOLABS<sup>®</sup> is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS