

NEBNext® dsDNA
Fragmentase®
Reaction Buffer v2



1-800-632-7799
info@neb.com
www.neb.com



B0349S 005160418041

B0349S

6 x 1.0 ml (6.0 ml) rxn buffer Lot: 0051604

Store at -20°C Exp: 4/18

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X NEBNext dsDNA Fragmentase

Reaction Buffer v2:

20 mM Tris-HCl
15 mM MgCl₂
50 mM NaCl
0.15% Triton X-100
0.1 mg/ml BSA
pH 7.5 @ 25°C

NEBNext® dsDNA
Fragmentase®
Reaction Buffer v2



1-800-632-7799
info@neb.com
www.neb.com



B0349S 005160418041

B0349S

6 x 1.0 ml (6.0 ml) rxn buffer Lot: 0051604

Store at -20°C Exp: 4/18

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X NEBNext dsDNA Fragmentase

Reaction Buffer v2:

20 mM Tris-HCl
15 mM MgCl₂
50 mM NaCl
0.15% Triton X-100
0.1 mg/ml BSA
pH 7.5 @ 25°C

Quality Control Assay

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

Protease Assay: Incubation of at least 1X NEBNext dsDNA Fragmentase Reaction Buffer v2 with 0.2 nmol of a standardized mixture of proteins, for 20 hours at 37°C, resulted in no proteolytic activity detected by SDS-PAGE.

Phosphatase Assay: Incubation of 10 µl of at least 1X NEBNext dsDNA Fragmentase Reaction Buffer v2 in protein phosphatase assay buffer (1 M diethanolamine @ pH 9.8 and 0.5 mM MgCl₂) containing 2.5 mM *p*-nitrophenyl phosphate at 37°C for 4 hours yields no detectable *p*-nitrophenylene anion as determined by spectrophotometric analysis at 405 nm.

Quality Control Assay

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

Protease Assay: Incubation of at least 1X NEBNext dsDNA Fragmentase Reaction Buffer v2 with 0.2 nmol of a standardized mixture of proteins, for 20 hours at 37°C, resulted in no proteolytic activity detected by SDS-PAGE.

Phosphatase Assay: Incubation of 10 µl of at least 1X NEBNext dsDNA Fragmentase Reaction Buffer v2 in protein phosphatase assay buffer (1 M diethanolamine @ pH 9.8 and 0.5 mM MgCl₂) containing 2.5 mM *p*-nitrophenyl phosphate at 37°C for 4 hours yields no detectable *p*-nitrophenylene anion as determined by spectrophotometric analysis at 405 nm.



NEW ENGLAND BIOLABS®, FRAGMENTASE® and NEBNEXT® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS®, FRAGMENTASE® and NEBNEXT® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS