

NiCo21(DE3) Competent *E. coli*



C2529H

20 x 0.05 ml/tube

Lot: 5

Store at -80°C

CAUTION: This product contains DMSO, a hazardous material. Review the MSDS before handling.

Description: Chemically competent *E. coli* cells derived from BL21(DE3). Poly-histidine tagged recombinant proteins that are isolated by immobilized metal affinity chromatography (IMAC) are often contaminated with significant amounts of endogenous *E. coli* metal binding proteins. The protein expression strain NiCo21(DE3) has been engineered to minimize *E. coli* protein contamination of IMAC fractions: GlnS is mutated to eliminate binding to IMAC resins and three other proteins (SlyD, ArnA and Can) are tagged to enable rapid removal by chitin affinity chromatography.

Features:

- Superior alternative to BL21(DE3) for routine protein expression
- Improved purity of target proteins isolated by IMAC
- Identical growth characteristics as BL21(DE3)
- Transformation efficiency: $1\text{--}5 \times 10^7$ cfu/ μg pUC19
- Deficient in proteases Lon and OmpT
- Resistant to phage T1 (*fhuA2*)
- Free of animal products

Reagents Supplied:

20 x 0.05 ml/tube of chemically competent NiCo21(DE3)
Competent *E. coli* cells (Store at -80°C)

20 ml of SOC Outgrowth Medium (Store at Room Temperature)

0.025 ml of 50 $\mu\text{g}/\text{ml}$ pUC19 Control DNA (Store at -20°C)

Compatible expression vectors/promoters:

T7 or T7-lac promoter, P_{lac} , P_{trc} , P_{lac} , Plac_{UV5} , ParaBAD vectors, PrhaBAD.

Note: Expression from vectors containing a T5-lacO promoter will not be regulated unless the expression vector also encodes the *lacI* gene or a compatible vector expresses *LacI*.

Quality Control Assays

Quality Control for Current Lot: Quality control values for a specific lot can be found on the datacard which accompanies each product.

Transformation Efficiency: 100 μg of pUC19 plasmid DNA was used to transform one tube of NiCo21(DE3) Competent *E. coli* following the protocol provided. $1\text{--}5 \times 10^7$ colonies formed/ μg after an overnight incubation on LB-ampicillin plates at 37°C .

Untransformed cells were also tested for resistance to phage $\phi 80$, a standard test for resistance to phage T1 and sensitivity to ampicillin, chloramphenicol, kanamycin, nitrofurantoin, spectinomycin, streptomycin and tetracycline.

STORAGE AND HANDLING: Competent cells should be stored at -80°C . Storage at -20°C will result in a significant decrease in transformation efficiency. Cells lose efficiency whenever they are warmed above -80°C , even if they do not thaw.



1-800-632-7799
info@neb.com
www.neb.com

High Efficiency Transformation Protocol

Perform steps 1–7 in the tube provided.

1. Thaw a tube of NiCo21(DE3) Competent *E. coli* cells on ice for 10 minutes.
2. Add 1–5 μl containing 1 μg –100 ng of plasmid DNA to the cell mixture. Carefully flick the tube 4–5 times to mix cells and DNA. **Do not vortex.**
3. Place the mixture on ice for 30 minutes. Do not mix.
4. Heat shock at exactly 42°C for exactly 10 seconds. Do not mix.
5. Place on ice for 5 minutes. Do not mix.
6. Pipette 950 μl of room temperature SOC into the mixture.
7. Place at 37°C for 60 minutes. Shake vigorously (250 rpm) or rotate.
8. Warm selection plates to 37°C .
9. Mix the cells thoroughly by flicking the tube and inverting, then perform several 10-fold serial dilutions in SOC.
10. Spread 50–100 μl of each dilution onto a selection plate and incubate overnight at 37°C . Alternatively, incubate at 30°C for 20–24 hours or at 25°C for 48 hours.

5 Minute Transformation Protocol

A shortened transformation protocol resulting in approximately 10% efficiency compared to the standard protocol may be suitable for applications where a reduced total number of transformants is acceptable. Follow the High Efficiency Transformation Protocol above with the following changes:

1. Steps 3 and 5 are reduced to 2 minutes.
2. Omit outgrowth (Step 7) completely for ampicillin-resistant plasmids or reduce the outgrowth time for other selective media as appropriate.

Protocol for Protein Expression Using NiCo21(DE3)

1. Transform expression plasmid into NiCo21(DE3). Plate on antibiotic selection plates and incubate overnight at 37°C .
2. Resuspend a single colony in liquid culture with antibiotic to produce a starter culture. Inoculate starter culture at a 1:100 dilution into expression media containing antibiotic.
3. Incubate at 37°C with shaking until OD_{600} reaches 0.4–0.8.
4. For most vector systems, induce expression with 40 or 400 μM IPTG for 3 hours at 37°C , 5 hours at 30°C or overnight at 16°C or 23°C .

Protocol for Removal of IMAC Contaminating Proteins:

E. coli SlyD, ArnA, and Can (carbonic anhydrase) are tagged with the chitin binding domain (CBD). Accordingly, these proteins may be removed by incubating the *E. coli* lysate or the IMAC elution fractions with chitin beads (NEB #S6651) or chitin magnetic beads (NEB #E8036). Binding of CBD-tagged proteins to chitin resin is compatible in a wide range of buffer conditions. Pooled IMAC fractions may be directly mixed with buffer-equilibrated chitin beads and incubated for 5–30 minutes to remove CBD-tagged contaminants from the His-tagged target protein.

The following procedure is recommended:

Use 1 ml of chitin resin for each volume of lysate or IMAC pool corresponding to 1 gram of NiCo21(DE3) cell pellet. (or use 1 ml of chitin resin for every 100 ml of expression culture). Resuspend chitin slurry (stored in 20% ethanol) and transfer to a gravity flow column. Equilibrate chitin column with buffer similar or equivalent to the IMAC low imidazole buffer: (or use a buffer compatible with the downstream chromatography step). Seal bottom of chitin column and add cell lysate or IMAC fractions containing CBD-tagged contaminants. Seal top of column and mix by rocking for 5–30 minutes at 4°C . Elute void volume containing target protein by gravity flow and optionally add extra equilibration buffer to displace all buffer containing the target protein.

(Alternatively, if using a mini-spin column incubate sample 5–30 minutes before centrifuging to elute the target protein).

Analyze eluted protein by SDS-PAGE or Western blot to determine purity. Removal of CBD-tagged contaminants may be analyzed by Anti-CBD Monoclonal Antibody (NEB #E8034).

(see other side)

Troubleshooting T7 Protein Expression

No colonies or no growth in liquid culture: This outcome is often due to basal expression of a target gene product, which is detrimental to cell viability. Basal expression of target protein in NiCo21 (DE3) is identical to basal expression in BL21 (DE3). Some vector systems (in particular those utilizing a T7 promoter) allow expression without inducer. If tightly regulated T7 expression is required, use a strain expressing *lysY*:

- T7 Express *lysY* (NEB #C3010) *lysY* produces a mutant T7 lysozyme which binds T7 RNA polymerase, reducing basal expression of the target protein. Upon induction, newly made T7 RNA polymerase titrates out the lysozyme and results in expression of the target protein.
- T7 Express *lysY/lq* (NEB #C3013) *lysY* expression as well as *lacI* over-expression to repress basal expression of the T7 RNA polymerase.
- Lemo21 (DE3) (NEB #C2528) BL21 (DE3) containing the Lemo System™. *LysY* expression is modulated by L-rhamnose, making T7 protein expression tightly regulated and tunable.

No protein visible by SDS-PAGE or no activity:

- Check for toxicity – no protein may mean the cells have lost the expression plasmid or elements of the expression plasmid have been deleted.
- Culture cells for protein induction. Just before induction, plate a sample on duplicate plates with and without antibiotic selection. If toxicity is an issue, there will be a significant difference between the number of colonies on the plates. Fewer colonies will be seen on plates containing antibiotic (indicating that the plasmid has been lost) compared to plates without antibiotic. If the expression plasmid is not maintained by drug selection, *lysY* strains will improve clone stability.
- Check clone integrity by restriction enzyme analysis and/or sequencing of the ORF.

Induced protein is insoluble: T7 expression often leads to very high production of protein that can result in the target protein becoming insoluble. Potential solutions are:

- Induce at a lower temperature (as low as 15°C overnight)
- Reduce IPTG concentration to 10–40 μ M
- Induce earlier in growth phase ($OD_{600} = 0.3$ or 0.4) and harvest cells earlier.
- Create an expression construct where the target protein is fused to Maltose Binding Protein (NEB #E8200).

Solutions/Recipes

SOB:		SOB:	
2%	Vegetable peptone (or Tryptone)	SOB + 20 mM Glucose	
0.5%	Yeast Extract		
10 mM	NaCl	LB agar:	
2.5 mM	KCl	1%	Tryptone
10 mM	MgCl ₂	0.5%	Yeast extract
10 mM	MgSO ₄	0.17 M	NaCl
		1.5%	Agar

Antibiotics for Plasmid Selection

Antibiotic	Working Concentration
Ampicillin	100 μ g/ml
Carbenicillin	100 μ g/ml
Chloramphenicol	33 μ g/ml
Kanamycin	30 μ g/ml
Streptomycin	25 μ g/ml
Tetracycline	15 μ g/ml

Genotype: *can::CBD fhuA2 [lon] ompT gal (λ DE3) [dcm] arnA::CBD slyD::CBD glmS6Ala Δ hds λ DE3 = λ sBamHlo Δ EcoRI-B int::(lacI::PlacUV5::T7 gene1) i21 Δ nin5*

Transformation Protocol Variables

Thawing: Cells are best thawed on ice and DNA added as soon as the last bit of ice in the tube disappears. Cells can also be thawed by hand, but warming above 0°C will decrease the transformation efficiency.

Incubation of DNA with Cells on Ice: For maximum transformation efficiency, cells and DNA should be incubated together on ice for 30 minutes. Expect a 2-fold loss in transformation efficiency for every 10 minutes you shorten this step.

Heat Shock: Both the temperature and the timing of the heat shock step are important and specific to the transformation volume and vessel. Using the transformation tube provided, 10 seconds at 42°C is optimal.

Outgrowth: Outgrowth at 37°C for 1 hour is best for cell recovery and for expression of antibiotic resistance. Expect a 2-fold loss in transformation efficiency for every 15 minutes you shorten this step. SOC gives 2-fold higher transformation efficiency than LB medium; and incubation with shaking or rotating the tube gives 2-fold higher transformation efficiency than incubation without shaking.

Plating: Selection plates can be used warm or cold, wet or dry without significantly affecting the transformation efficiency. However, warm, dry plates are easier to spread and allow for the most rapid colony formation.

Strain Properties

The properties of this strain that contribute to its usefulness as a protein expression strain are described below.

T7 RNA Polymerase: *T7 gene1* is encoded by the lambda DE3 prophage present within the chromosome. T7 RNA polymerase is expressed from the *lacUV5* promoter, which is less sensitive to catabolite repression than the *wt lac* promoter. Thus DE3 strains may exhibit uninduced target protein expression. Although λ DE3 is normally dormant in the host chromosome, the induction of the SOS response can occur as the result of expressing proteins that damage the *E. coli* chromosome, either directly or indirectly. This may lead to cell lysis. T7 Express strains do not carry the DE3 prophage and better tolerate an SOS response.

Protease Deficient ([*lon ompT*): *E. coli* B strains are “naturally” deficient in the Lon protease which in K-12 strains serves to degrade misfolded proteins and to prevent some cell cycle-specific proteins from accumulating. The OmpT protease resides at the surface of wild type *E. coli* in both K-12 and B strains, presumably helping the cells to derive amino acids from their external environment. Cells deficient in both these proteases are much more amenable to the production of proteins from cloned genes.

T1 Phage Resistant (*fhuA2*): T1, an extremely virulent phage requires the *E. coli* ferric hydroxamate uptake receptor for infectivity. Deletion of this gene confers resistance to this type of phage, but does not significantly affect the transformation or growth characteristics of the cell.

***glmS6Ala*:** The *glmS* gene (glucosamine synthetase) is mutated and the expressed GlmS protein (67 kDa) contains six histidine to alanine substitutions (positions 62, 65, 432, 436, 466, 467). The mutated GlmS protein does not bind Ni-NTA resin in the presence of 20mM imidazole binding/wash buffer, whereas wt GlmS protein binds Ni-NTA resin and is not eluted until the imidazole concentration is within 55–80 mM (1).

arnA::CBD The *arnA* gene (lipid A modification) is fused with a *B. circulans* ORF encoding a chitin binding domain. The expressed fusion protein is 82 kDa whereas the wt ArnA protein is 74 kDa.

can::CBD The *can* gene (carbonic anhydrase) is fused with a *B. circulans* ORF encoding a chitin binding domain. The expressed fusion protein is 32 kDa whereas wt carbonic anhydrase is 25 kDa.

slyD::CBD The *slyD* gene (FKBP-type prolyl isomerase) is fused with a *B. circulans* ORF encoding a chitin binding domain. The expressed fusion protein is 28 kDa. However, the apparent mass of the CBD fusion is approximately 35 kDa when analyzed by SDS-PAGE. The calculated mass of wt SlyD is 21 kDa, whereas the apparent mass is 28 kDa.

The GlmS(6Ala) protein and the CBD-tagged proteins are functional *in vivo* according to assays conducted at New England Biolabs.

Reference:

1. Bolanos-Garcia and Davies (2006) *Biochimica et Biophysica Acta* 1760, 1304–1313.

Notice to Buyer/User: The buyer/user has a non-exclusive license to use this system or any components thereof for Research Purposes Only. **RESEARCH USE ASSURANCE STATEMENT** The buyer and user have a non-exclusive sub-license to use this system or any component thereof for RESEARCH PURPOSES ONLY, based upon agreement to the following assurances: Transfer of the host cells that contain the cloned copy of the T7 gene 1 to third parties is explicitly prohibited. This information applies to *E. coli* ER2566, ER2833, ER3011, ER3012, ER3013 and ER3021, SHuffle™ T7, SHuffle™ T7 LysY, SHuffle™ T7 Express, SHuffle™ T7 Express LysY and their competent derivatives, C2566, C2833, C3010, C3013, C3016, C3022, C3026, C3027, C3029 and C3030 when provided separately or when provided in combination with appropriate vectors for said systems. A license to use this system or any components thereof for commercial purposes may be obtained from New England Biolabs, Inc. **Commercial Laboratory Buyer and User:** Use of the host cells ER2566, ER2833, C2566, C2833, C3010, C3013, C3016, C3022, C3026, C3027, C3029, C3030, SHuffle™ T7, SHuffle™ T7 LysY, SHuffle™ T7 Express, SHuffle™ T7 Express LysY or their competent derivatives that contain the cloned copy of T7 gene 1, the gene for T7 RNA polymerase for any purpose other than in combination with either a T7/pMAL or T7/IMPACT vector is explicitly prohibited. Use of the host cells that may contain the cloned copy of the T7 gene 1, the gene for T7 RNA polymerase with any other vector(s) containing a T7 promoter to direct the production of RNA or protein requires a license from Brookhaven National Laboratory. Information about research-use or commercial-use license agreement may be obtained from the Office of Intellectual Property and Sponsored Research, Brookhaven National Laboratory, Building 475D, P.O. Box 5000, Upton, New York 11973-5000. Tel.: 631-344-7134. Fax: 631-344-3729. You may refuse this non-exclusive research license agreement by returning the enclosed materials unused. By keeping or using the enclosed materials, you agree to be bound by the terms of this sub-license. **ACADEMIC AND NON-PROFIT LABORATORY ASSURANCE LETTER** The T7 expression system is based on technology developed at Brookhaven National Laboratory under contract with the U.S. Department of Energy and is the subject of patent applications assigned to Brookhaven Science Associates, LLC (BSA). BSA will grant a non-exclusive license for use of this technology, including the enclosed materials, based upon the following assurances: 1. These materials are to be used for non-commercial research purposes only. A separate license is required for any commercial use, including the use of these materials for research purposes or production purposes by any commercial entity. Information about commercial licenses may be obtained from the Office of Intellectual Property and Sponsored Research, Brookhaven National Laboratory, Building 475D, P.O. Box 5000, Upton, New York 11973-5000. Tel: 631-344-7134. 2. No materials that contain the cloned copy of T7 gene 1, the gene for T7 RNA polymerase, may be distributed further to third parties outside of your laboratory, unless recipient receives a copy of this license and agrees to be bound by the terms. This limitation applies to strains ER2566, ER2833, C2566, C2833, C3010, C3013, C3016, C3022, C3026, C3027, C3029, C3030, BL21 (DE3), BL21 (DE3)pLysS and BL21 (DE3)pLysE, SHuffle™ T7, SHuffle™ T7 LysY, SHuffle™ T7 Express, SHuffle™ T7 Express LysY and other competent derivatives and any derivatives you may make of them, including such strains containing recombinant vectors. You may refuse this license by returning the enclosed materials unused. By keeping or using the enclosed materials, you agree to be bound by the terms of this license.