

Taq DNA Ligase



1-800-632-7799
info@neb.com
www.neb.com



M0208S 011160818081

M0208S



2,000 units **40,000 U/ml** **Lot: 0111608**

RECOMBINANT **Store at -20°C** **Exp: 8/18**

Description: *Taq* DNA Ligase catalyzes the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini of two adjacent oligonucleotides which are hybridized to a complementary target DNA. The ligation will occur only if the oligonucleotides are perfectly paired to the complementary target DNA and have no gaps between them; therefore, a single-base substitution can be detected. *Taq* DNA Ligase is active at elevated temperatures (45–65°C) (1,2).

Taq DNA Ligase



1-800-632-7799
info@neb.com
www.neb.com



M0208S 011160818081

M0208S



2,000 units **40,000 U/ml** **Lot: 0111608**

RECOMBINANT **Store at -20°C** **Exp: 8/18**

Description: *Taq* DNA Ligase catalyzes the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini of two adjacent oligonucleotides which are hybridized to a complementary target DNA. The ligation will occur only if the oligonucleotides are perfectly paired to the complementary target DNA and have no gaps between them; therefore, a single-base substitution can be detected. *Taq* DNA Ligase is active at elevated temperatures (45–65°C) (1,2).

Source: Purified from an *E. coli* strain containing the cloned ligase gene from *Thermus aquaticus* HB8 (1)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Applications:

- Allele-specific gene detection using Ligase Detection Reaction and Ligase Chain Reaction (1,3).
- Mutagenesis by incorporation of a phosphorylated oligonucleotide during PCR amplification (4).

Reagents Supplied with Enzyme: 10X *Taq* DNA Ligase Reaction Buffer.

Reaction Conditions: Incubate DNA and enzyme in 1X *Taq* DNA Ligase Buffer at 45°C for 15 minutes or in a thermocycler with a program suited to the reaction described by Barany (1991) Genetic Disease Detection and DNA Amplification Using Cloned Thermostable Ligase. *Proc. Natl. Acad. Sci. USA* 88, 189–193. "The reaction is stopped with a mixture of 50% glycerol, 50 mM EDTA, bromophenol blue."

Source: Purified from an *E. coli* strain containing the cloned ligase gene from *Thermus aquaticus* HB8 (1)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Applications:

- Allele-specific gene detection using Ligase Detection Reaction and Ligase Chain Reaction (1,3).
- Mutagenesis by incorporation of a phosphorylated oligonucleotide during PCR amplification (4).

Reagents Supplied with Enzyme: 10X *Taq* DNA Ligase Reaction Buffer.

Reaction Conditions: Incubate DNA and enzyme in 1X *Taq* DNA Ligase Buffer at 45°C for 15 minutes or in a thermocycler with a program suited to the reaction described by Barany (1991) Genetic Disease Detection and DNA Amplification Using Cloned Thermostable Ligase. *Proc. Natl. Acad. Sci. USA* 88, 189–193. "The reaction is stopped with a mixture of 50% glycerol, 50 mM EDTA, bromophenol blue."

1X *Taq* DNA Ligase Reaction Buffer:

20 mM Tris-HCl
25 mM potassium acetate
10 mM magnesium acetate
10 mM dithiothreitol
1 mM NAD
0.1% Triton X-100
(pH 7.6 @ 25°C)

Requires NAD⁺ as a cofactor. NAD⁺ is supplied in the 10X *Taq* DNA Ligase Reaction Buffer; the buffer should be stored at -70°C to extend the half life of the NAD⁺ cofactor.

Unit Definition: (Cohesive End Unit)

One unit is defined as the amount of enzyme required to give 50% ligation of the 12-base pair cohesive ends of 1 µg of BstEII-digested λ DNA in a total reaction volume of 50 µl in 15 minutes at 45°C.

Unit Assay Conditions: 1X *Taq* DNA Ligase Reaction Buffer and DNA (20 µg/ml). After incubation at 45°C for 15 minutes, the reaction is terminated by addition of stop dye (50% glycerol, 50 mM EDTA and bromophenol blue), heated at 70°C for 10 minutes and then loaded on a 0.7% agarose gel. Due to the presence of ligase, the cos ends of BstEII-digested λ DNA will stay together after 70°C heat treatment.

1X *Taq* DNA Ligase Reaction Buffer:

20 mM Tris-HCl
25 mM potassium acetate
10 mM magnesium acetate
10 mM dithiothreitol
1 mM NAD
0.1% Triton X-100
(pH 7.6 @ 25°C)

Requires NAD⁺ as a cofactor. NAD⁺ is supplied in the 10X *Taq* DNA Ligase Reaction Buffer; the buffer should be stored at -70°C to extend the half life of the NAD⁺ cofactor.

Unit Definition: (Cohesive End Unit)

One unit is defined as the amount of enzyme required to give 50% ligation of the 12-base pair cohesive ends of 1 µg of BstEII-digested λ DNA in a total reaction volume of 50 µl in 15 minutes at 45°C.

Unit Assay Conditions: 1X *Taq* DNA Ligase Reaction Buffer and DNA (20 µg/ml). After incubation at 45°C for 15 minutes, the reaction is terminated by addition of stop dye (50% glycerol, 50 mM EDTA and bromophenol blue), heated at 70°C for 10 minutes and then loaded on a 0.7% agarose gel. Due to the presence of ligase, the cos ends of BstEII-digested λ DNA will stay together after 70°C heat treatment.

Heat Inactivation: No

Quality Control Assays

Exonuclease Activity: Incubation of 1,200 units for 4 hours at 37°C in 50 µl of thermostable ligase buffer containing 1 µg sonicated ³H DNA (10⁵ cpm/µg) gave < 0.1% acid soluble counts.

Non-Specific DNase Activity (16 hours): A 50 µl reaction in NEBuffer 4 containing 1 µg of λ HindIII DNA and 80 units of *Taq* DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Endonuclease Activity: Incubation of 1,500 units of enzyme for 4 hours at 37°C in 50 µl of assay buffer without NAD and 1 µg φX174 RF I DNA gave < 10% conversion to RF II.

(see other side)

CERTIFICATE OF ANALYSIS

Heat Inactivation: No

Quality Control Assays

Exonuclease Activity: Incubation of 1,200 units for 4 hours at 37°C in 50 µl of thermostable ligase buffer containing 1 µg sonicated ³H DNA (10⁵ cpm/µg) gave < 0.1% acid soluble counts.

Non-Specific DNase Activity (16 hours): A 50 µl reaction in NEBuffer 4 containing 1 µg of λ HindIII DNA and 80 units of *Taq* DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Endonuclease Activity: Incubation of 1,500 units of enzyme for 4 hours at 37°C in 50 µl of assay buffer without NAD and 1 µg φX174 RF I DNA gave < 10% conversion to RF II.

(see other side)

CERTIFICATE OF ANALYSIS

References:

1. Barany, F. (1991) *Proc. Natl. Acad. Sci. USA* 88, 189-193.
2. Takahashi, M. et al. (1984) *J. Biol. Chem.* 259, 10041–10047.
3. Barany, F. (1991) *The Ligase Chain Reaction in a PCR World* (pp. 5–16). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
4. Michael, S.F. (1994) *Biotechniques* 16, 411–412.

Page 2 (M0208)

References:

1. Barany, F. (1991) *Proc. Natl. Acad. Sci. USA* 88, 189-193.
2. Takahashi, M. et al. (1984) *J. Biol. Chem.* 259, 10041–10047.
3. Barany, F. (1991) *The Ligase Chain Reaction in a PCR World* (pp. 5–16). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
4. Michael, S.F. (1994) *Biotechniques* 16, 411–412.

Page 2 (M0208)

V1.0

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



This product is covered by one or more patents, trademarks and/or copyrights owned or controlled by New England Biolabs, Inc. For more information, please contact NEB's Global Business Development team at gbd@neb.com.

Taq DNA Ligase: Notice to Purchaser: This product is designed to ligate DNA fragments at temperatures requiring a thermoactive and thermostable enzyme. The seller is aware that the product may be used in the Ligase Chain Reaction™ (LCR™) process covered by one of more claims of a pending patent application or issued patent assigned to Cornell Research Foundation Inc., or Cornell Research Foundation, Inc. and the California Institute of Technology. LCR™ license inquires should be directed to Cornell Research Foundation, Inc.

V1.0

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



This product is covered by one or more patents, trademarks and/or copyrights owned or controlled by New England Biolabs, Inc. For more information, please contact NEB's Global Business Development team at gbd@neb.com.

Taq DNA Ligase: Notice to Purchaser: This product is designed to ligate DNA fragments at temperatures requiring a thermoactive and thermostable enzyme. The seller is aware that the product may be used in the Ligase Chain Reaction™ (LCR™) process covered by one of more claims of a pending patent application or issued patent assigned to Cornell Research Foundation Inc., or Cornell Research Foundation, Inc. and the California Institute of Technology. LCR™ license inquires should be directed to Cornell Research Foundation, Inc.