

## DNA Polymerase I, Large (Klenow) Fragment



1-800-632-7799  
info@neb.com  
www.neb.com



M0210S 089140316031

# M0210S



**200 units**      **5,000 U/ml**      **Lot: 0891403**  
**RECOMBINANT**    **Store at -20°C**    **Exp: 3/16**

**Description:** DNA Polymerase I, Large (Klenow) Fragment is a proteolytic product of *E. coli* DNA Polymerase I which retains polymerization and 3'→5' exonuclease activity, but has lost 5'→3' exonuclease activity (1). Klenow retains the polymerization fidelity of the holoenzyme without degrading 5' termini.

**Source:** Purified from a strain of *E. coli* that carries the DNA Polymerase I, Large (Klenow) Fragment gene.

### Applications:

- DNA sequencing by the Sanger dideoxy method (2)
- Fill-in of 5' overhangs to form blunt ends (3)
- Removal of 3' overhangs to form blunt ends (3)
- Second strand cDNA synthesis
- Second strand synthesis in mutagenesis protocols (4)

Supplied in: 25 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol and 50% glycerol.

**Reagents Supplied with Enzyme:**  
10X NEBuffer 2

**Reaction Conditions:** 1X NEBuffer 2.  
Supplement with dNTPs (not included).

Klenow Fragment is also active in all four NEBuffers and T4 DNA Ligase Reaction Buffer when supplemented with dNTPs.

### 1X NEBuffer 2:

50 mM NaCl  
10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>  
1 mM DTT  
pH 7.9 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 37°C.

**Unit Assay Conditions:** 1X NEBuffer 2, 33 μM dNTPs including [<sup>3</sup>H]-dTTP and 70 μg/ml denatured herring sperm DNA.

**DNA Sequencing:** When this preparation is used to sequence DNA using the dideoxy method of Sanger et al., 1 unit/5 μl reaction volume is recommended.

**Molecular Weight:** 68,000 daltons.

**Heat Inactivation:** 75°C for 20 minutes.

### Quality Control Assays

**Endonuclease Activity:** Incubation of a 50 μl reaction in NEBuffer 2 containing a minimum of 50 units of DNA Polymerase I, Large (Klenow) Fragment with 1 μg of supercoiled φX174 DNA for 4 hours at 37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Notes on Use:** Protocol for blunting ends by 3' overhang removal and fill-in of 3' recessed (5' overhang) end: DNA should be dissolved in 1X NEBuffer 1-4 or T4 DNA Ligase Reaction Buffer supplemented with 33 μM each dNTP. Add 1 unit Klenow per microgram DNA and incubate 15 minutes at 25°C. Stop reaction by adding EDTA to a final concentration of 10 mM and heating at 75°C for 20 minutes.

**CAUTION:** Elevated temperatures, excessive amounts of enzyme, failure to supplement with dNTPs or long reaction times will result in recessed ends due to the 3'→5' exonuclease activity of the enzyme.

(see other side)

CERTIFICATE OF ANALYSIS

## DNA Polymerase I, Large (Klenow) Fragment



1-800-632-7799  
info@neb.com  
www.neb.com



M0210S 089140316031

# M0210S



**200 units**      **5,000 U/ml**      **Lot: 0891403**  
**RECOMBINANT**    **Store at -20°C**    **Exp: 3/16**

**Description:** DNA Polymerase I, Large (Klenow) Fragment is a proteolytic product of *E. coli* DNA Polymerase I which retains polymerization and 3'→5' exonuclease activity, but has lost 5'→3' exonuclease activity (1). Klenow retains the polymerization fidelity of the holoenzyme without degrading 5' termini.

**Source:** Purified from a strain of *E. coli* that carries the DNA Polymerase I, Large (Klenow) Fragment gene.

### Applications:

- DNA sequencing by the Sanger dideoxy method (2)
- Fill-in of 5' overhangs to form blunt ends (3)
- Removal of 3' overhangs to form blunt ends (3)
- Second strand cDNA synthesis
- Second strand synthesis in mutagenesis protocols (4)

Supplied in: 25 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol and 50% glycerol.

**Reagents Supplied with Enzyme:**  
10X NEBuffer 2

**Reaction Conditions:** 1X NEBuffer 2.  
Supplement with dNTPs (not included).

Klenow Fragment is also active in all four NEBuffers and T4 DNA Ligase Reaction Buffer when supplemented with dNTPs.

### 1X NEBuffer 2:

50 mM NaCl  
10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>  
1 mM DTT  
pH 7.9 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 37°C.

**Unit Assay Conditions:** 1X NEBuffer 2, 33 μM dNTPs including [<sup>3</sup>H]-dTTP and 70 μg/ml denatured herring sperm DNA.

**DNA Sequencing:** When this preparation is used to sequence DNA using the dideoxy method of Sanger et al., 1 unit/5 μl reaction volume is recommended.

**Molecular Weight:** 68,000 daltons.

**Heat Inactivation:** 75°C for 20 minutes.

### Quality Control Assays

**Endonuclease Activity:** Incubation of a 50 μl reaction in NEBuffer 2 containing a minimum of 50 units of DNA Polymerase I, Large (Klenow) Fragment with 1 μg of supercoiled φX174 DNA for 4 hours at 37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Notes on Use:** Protocol for blunting ends by 3' overhang removal and fill-in of 3' recessed (5' overhang) end: DNA should be dissolved in 1X NEBuffer 1-4 or T4 DNA Ligase Reaction Buffer supplemented with 33 μM each dNTP. Add 1 unit Klenow per microgram DNA and incubate 15 minutes at 25°C. Stop reaction by adding EDTA to a final concentration of 10 mM and heating at 75°C for 20 minutes.

**CAUTION:** Elevated temperatures, excessive amounts of enzyme, failure to supplement with dNTPs or long reaction times will result in recessed ends due to the 3'→5' exonuclease activity of the enzyme.

(see other side)

CERTIFICATE OF ANALYSIS

**References:**

1. Jacobsen, H., Klenow, H. and Overgaard-Hansen, K. (1974) *Eur. J. Biochem.* 45, 623–627.
2. Sanger, F. et al. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
3. Sambrook, J. et al. (1989). *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (pp. 5.40–5.43). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
4. Gubler, U. (1987). In S.L. Berger and A.R. Kimmel (Eds.), *Methods in Enzymology*, Vol.152, (pp. 330–335). San Diego: Academic Press.

**Companion Products Sold Separately:**

NEBuffer 2  
#B7002S                      6.0 ml

Deoxynucleotide Solution Set  
#N0446S      25 µmol of each

Deoxynucleotide Solution Mix  
#N0447S      8 µmol of each  
#N0447L      40 µmol of each

**References:**

1. Jacobsen, H., Klenow, H. and Overgaard-Hansen, K. (1974) *Eur. J. Biochem.* 45, 623–627.
2. Sanger, F. et al. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463–5467.
3. Sambrook, J. et al. (1989). *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (pp. 5.40–5.43). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
4. Gubler, U. (1987). In S.L. Berger and A.R. Kimmel (Eds.), *Methods in Enzymology*, Vol.152, (pp. 330–335). San Diego: Academic Press.

**Companion Products Sold Separately:**

NEBuffer 2  
#B7002S                      6.0 ml

Deoxynucleotide Solution Set  
#N0446S      25 µmol of each

Deoxynucleotide Solution Mix  
#N0447S      8 µmol of each  
#N0447L      40 µmol of each