

## Deoxynucleotide (dNTP) Solution Set



1-800-632-7799  
info@neb.com  
www.neb.com



N0446S 053150217022

# N0446S

25 µmol each dNTP

Lot: 0531502

Store at -20°C

Exp: 2/17

**Description:** The Deoxynucleotide (dNTP) Solution Set contains 0.25 ml of each 100 mM ultrapure deoxynucleotide as four separate solutions (dATP, dCTP, dGTP and dTTP).

Supplied in: Ultrapure water as a sodium salt at pH 7.5.

**Concentration:** Each deoxynucleotide is supplied at a concentration of 100 mM ( $\pm 5$  mM) as determined by  $A_{260}$  absorbance.

### Reagents Supplied:

100 mM dATP Solution	0.25 ml
100 mM dCTP Solution	0.25 ml
100 mM dGTP Solution	0.25 ml
100 mM dTTP Solution	0.25 ml

**Diluent Compatibility:** Can be diluted using sterile distilled water, preferably Milli-Q® water or can be diluted using sterile TE [10 mM Tris-HCl, 1 mM EDTA (pH 7.5)].

### Quality Controls

The purity of each deoxynucleotide is  $\geq 99\%$  as determined by HPLC analysis.

**0.5 kb, 2 kb and 5 kb Lambda PCR Assay –** 25 cycles of PCR amplification of 1 ng Lambda DNA with 5 units of *Taq* DNA Polymerase in the presence of 200 µM dATP, dCTP, dGTP and dTTP 0.5 µM primers and 1X ThermoPol® Buffer results in the amplification of the specific 0.5 kb, 2 kb and 5 kb products as determined by agarose gel electrophoresis.

### Phosphatase Activity Assay (pNPP Colorimetric Assay) –

A protein phosphatase buffer solution containing 2 mM dATP, dCTP, dGTP or dTTP solution and 100 µM *p*-nitrophenol phosphate, incubated for 4 hours at 37°C, yields no detectable phosphatase activity as determined by spectrophotometric analysis of released *p*-nitrophenylene anion at 405 nm.

**Non-specific Nuclease Assay –** A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of T3 DNA or HindIII digested Lambda DNA and a minimum of 10 µl of dATP, dCTP, dGTP or dTTP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

For use with any DNA Polymerase from NEB.

### Companion Products Sold Separately:

Deoxynucleotide (dNTP) Solution Mix  
#N0447S 8 µmol of each  
#N0447L 40 µmol of each

dATP Solution  
#N0440S 25 µmol



NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

MILLI-Q® is a registered trademark of Millipore Corporation.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

## Deoxynucleotide (dNTP) Solution Set



1-800-632-7799  
info@neb.com  
www.neb.com



N0446S 053150217022

# N0446S

25 µmol each dNTP

Lot: 0531502

Store at -20°C

Exp: 2/17

**Description:** The Deoxynucleotide (dNTP) Solution Set contains 0.25 ml of each 100 mM ultrapure deoxynucleotide as four separate solutions (dATP, dCTP, dGTP and dTTP).

Supplied in: Ultrapure water as a sodium salt at pH 7.5.

**Concentration:** Each deoxynucleotide is supplied at a concentration of 100 mM ( $\pm 5$  mM) as determined by  $A_{260}$  absorbance.

### Reagents Supplied:

100 mM dATP Solution	0.25 ml
100 mM dCTP Solution	0.25 ml
100 mM dGTP Solution	0.25 ml
100 mM dTTP Solution	0.25 ml

**Diluent Compatibility:** Can be diluted using sterile distilled water, preferably Milli-Q® water or can be diluted using sterile TE [10 mM Tris-HCl, 1 mM EDTA (pH 7.5)].

### Quality Controls

The purity of each deoxynucleotide is  $\geq 99\%$  as determined by HPLC analysis.

**0.5 kb, 2 kb and 5 kb Lambda PCR Assay –** 25 cycles of PCR amplification of 1 ng Lambda DNA with 5 units of *Taq* DNA Polymerase in the presence of 200 µM dATP, dCTP, dGTP and dTTP 0.5 µM primers and 1X ThermoPol® Buffer results in the amplification of the specific 0.5 kb, 2 kb and 5 kb products as determined by agarose gel electrophoresis.

### Phosphatase Activity Assay (pNPP Colorimetric Assay) –

A protein phosphatase buffer solution containing 2 mM dATP, dCTP, dGTP or dTTP solution and 100 µM *p*-nitrophenol phosphate, incubated for 4 hours at 37°C, yields no detectable phosphatase activity as determined by spectrophotometric analysis of released *p*-nitrophenylene anion at 405 nm.

**Non-specific Nuclease Assay –** A 50 µl reaction in 1X NEBuffer 2 containing 1 µg of T3 DNA or HindIII digested Lambda DNA and a minimum of 10 µl of dATP, dCTP, dGTP or dTTP Solution incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

For use with any DNA Polymerase from NEB.

### Companion Products Sold Separately:

Deoxynucleotide (dNTP) Solution Mix  
#N0447S 8 µmol of each  
#N0447L 40 µmol of each

dATP Solution  
#N0440S 25 µmol



NEW ENGLAND BIOLABS® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

MILLI-Q® is a registered trademark of Millipore Corporation.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS