

α 1-3, 6 Galactosidase



1-800-632-7799
info@neb.com
www.neb.com



P0731S 004150816081

P0731S




100 units 4,000 U/ml Lot: 0041508

RECOMBINANT Store at 4°C Exp: 8/16

Description: α 1-3, 6 Galactosidase is a highly specific exoglycosidase that catalyzes the hydrolysis of α 1-3, 6 linked D-galactopyranosyl residues from oligosaccharides.

Specificity:



α 1-3, 6 Galactosidase



1-800-632-7799
info@neb.com
www.neb.com



P0731S 004150816081

P0731S




100 units 4,000 U/ml Lot: 0041508

RECOMBINANT Store at 4°C Exp: 8/16


Description: α 1-3, 6 Galactosidase is a highly specific exoglycosidase that catalyzes the hydrolysis of α 1-3, 6 linked D-galactopyranosyl residues from oligosaccharides.

Specificity:




Detailed Specificity: Specificity can vary depending on incubation time and branching structure.

A) 0.1 nm/ μ l substrate, 1 hour incubation



B) 0.1 nm/ μ l substrate, 18 hour incubation



C) 0.1 nm/ μ l substrate, 1 hour incubation, not cleaved






Figure 1: Detailed specificity of α 1-3, 6 Galactosidase. Reactions (A) and (C) contained 4 units of α 1-3, 6 Galactosidase, 1X GlycoBuffer 1 and 1X BSA in a total reaction volume of 10 μ l. Reaction (C) shows that branched fucose inhibits cleavage. Reaction (B) contained 24 units of α 1-3, 6 Galactosidase and 100 units of Neuraminidase, followed by a heat kill at 65°C for 10 minutes and a 2 hour digestion with 16 units of β 1-4 Galactosidase. The reaction in (B) contained 1X GlycoBuffer 1 and 1X BSA in a total reaction volume of 20 μ l. The reactions were incubated at 37°C. Complete digestion of the α 1-3, 6 Galactosidase was determined by an observation of complete transformation of the substrate in (B) to the non-reducing terminal N-acetylglucosamine tetra antennary oligosaccharide.

Detailed Specificity: Specificity can vary depending on incubation time and branching structure.

A) 0.1 nm/ μ l substrate, 1 hour incubation



B) 0.1 nm/ μ l substrate, 18 hour incubation



C) 0.1 nm/ μ l substrate, 1 hour incubation, not cleaved




Figure 1: Detailed specificity of α 1-3, 6 Galactosidase. Reactions (A) and (C) contained 4 units of α 1-3, 6 Galactosidase, 1X GlycoBuffer 1 and 1X BSA in a total reaction volume of 10 μ l. Reaction (C) shows that branched fucose inhibits cleavage. Reaction (B) contained 24 units of α 1-3, 6 Galactosidase and 100 units of Neuraminidase, followed by a heat kill at 65°C for 10 minutes and a 2 hour digestion with 16 units of β 1-4 Galactosidase. The reaction in (B) contained 1X GlycoBuffer 1 and 1X BSA in a total reaction volume of 20 μ l. The reactions were incubated at 37°C. Complete digestion of the α 1-3, 6 Galactosidase was determined by an observation of complete transformation of the substrate in (B) to the non-reducing terminal N-acetylglucosamine tetra antennary oligosaccharide.

Source: Cloned from *Xanthomonas manihotis* and expressed in *E. coli* (1).

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na₂EDTA.

Reagents Supplied with Enzyme:

10X GlycoBuffer 1
100X BSA

Reaction Conditions:

1X GlycoBuffer 1:
50 mM Sodium Acetate (pH 5.5 @ 25°C)
and 5 mM CaCl₂. Supplement with 100 μ g/ml BSA.
Incubate at 37°C.

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the terminal, α -D-galactose from 1 nmol Gal α 1-3Gal β 1-4Gal-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.

(See other side)

CERTIFICATE OF ANALYSIS

Source: Cloned from *Xanthomonas manihotis* and expressed in *E. coli* (1).

Supplied in: 50 mM NaCl, 20 mM Tris-HCl (pH 7.5 @ 25°C) and 1 mM Na₂EDTA.

Reagents Supplied with Enzyme:

10X GlycoBuffer 1
100X BSA

Reaction Conditions:

1X GlycoBuffer 1:
50 mM Sodium Acetate (pH 5.5 @ 25°C)
and 5 mM CaCl₂. Supplement with 100 μ g/ml BSA.
Incubate at 37°C.

Optimal incubation times and enzyme concentrations must be determined empirically for a particular substrate.

Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the terminal, α -D-galactose from 1 nmol Gal α 1-3Gal β 1-4Gal-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.

(See other side)

CERTIFICATE OF ANALYSIS

Unit Definition Assay: Two fold serial dilutions of α 1-3, 6 Galactosidase are incubated with 1 nmol AMC-labeled substrate in 1X GlycoBuffer 1 and 1X BSA in a 10 μ l reaction. The reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized via thin layer chromatography (2).

Specific Activity: 137,000 units/mg

Molecular Weight: 70,000 daltons.

Quality Assurance: No contaminating exoglycosidase or proteolytic activity could be detected.

Quality Controls

Glycosidase Assays:

12 units of α 1-3, 6 Galactosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 μ l reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

Page 2 (P0731)

Unit Definition Assay: Two fold serial dilutions of α 1-3, 6 Galactosidase are incubated with 1 nmol AMC-labeled substrate in 1X GlycoBuffer 1 and 1X BSA in a 10 μ l reaction. The reaction mix is incubated for 1 hour at 37°C. Separation of reaction products are visualized via thin layer chromatography (2).

Specific Activity: 137,000 units/mg

Molecular Weight: 70,000 daltons.

Quality Assurance: No contaminating exoglycosidase or proteolytic activity could be detected.

Quality Controls

Glycosidase Assays:

12 units of α 1-3, 6 Galactosidase were incubated with 0.1 mM of fluorescently-labeled oligosaccharides and glycopeptides, in a 10 μ l reaction for 20 hours at 37°C. The reaction products were analyzed by TLC for digestion of substrate.

Page 2 (P0731)

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

No other glycosidase activities were detected (ND) with the following substrates:

β -N-Acetylglucosaminidase:
GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC ND

α -N-Acetylgalactosaminidase:
GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC ND

α -Fucosidase:
Fuc α 1-2Gal β 1-4Glc-AMC ND

Gal β 1-4 (Fuc α 1-3)GlcNAc β 1-3Gal β 1-4Glc-AMC ND

β -Galactosidase:
Gal β 1-3GlcNAc β 1-4Gal β 1-4Glc-AMC ND

Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC ND

α -Neuraminidase:
Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC ND

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

No other glycosidase activities were detected (ND) with the following substrates:

β -N-Acetylglucosaminidase:
GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC ND

α -N-Acetylgalactosaminidase:
GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC ND

α -Fucosidase:
Fuc α 1-2Gal β 1-4Glc-AMC ND

Gal β 1-4 (Fuc α 1-3)GlcNAc β 1-3Gal β 1-4Glc-AMC ND

β -Galactosidase:
Gal β 1-3GlcNAc β 1-4Gal β 1-4Glc-AMC ND

Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC ND

α -Neuraminidase:
Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC ND

α -Mannosidase:
Man α 1-3Man β 1-4GlcNAc-AMC ND

Man α 1-6Man α 1-6(Man α 1-3)Man-AMC ND

β -Glucosidase:
Glc β 1-4Glc β 1-4Glc-AMC ND

α -Glucosidase:
Glc α 1-6Glc α 1-4Glc-AMC ND

β -Xylosidase:
Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC ND

β -Mannosidase:
Man β 1-4Man β 1-4Man-AMC ND

Endo F₁, F₂, H:
Dansylated invertase high mannose. ND

Endo F₂, F₃:
Dansylated fibrinogen biantennary. ND

PNGase F:
Fluoresceinated fetuin triantennary. ND

α -Mannosidase:
Man α 1-3Man β 1-4GlcNAc-AMC ND

Man α 1-6Man α 1-6(Man α 1-3)Man-AMC ND

β -Glucosidase:
Glc β 1-4Glc β 1-4Glc-AMC ND

α -Glucosidase:
Glc α 1-6Glc α 1-4Glc-AMC ND

β -Xylosidase:
Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC ND

β -Mannosidase:
Man β 1-4Man β 1-4Man-AMC ND

Endo F₁, F₂, H:
Dansylated invertase high mannose. ND

Endo F₂, F₃:
Dansylated fibrinogen biantennary. ND

PNGase F:
Fluoresceinated fetuin triantennary. ND

Protease Assay: After incubation of 28 units of α 1-3, 6 Galactosidase with 0.2 nmol of a standard mixture of proteins in a 20 μ l reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

Note: Recommended storage temperature is 4°C. Avoid repeated freeze/thaw cycles

Heat Inactivation: 65°C for 10 minutes.

References:

1. McLeod, E., New England Biolabs, Inc. unpublished results.
2. Wong-Madden, S.T. and Landry, D. (1995) *Glycobiology* 5, 19–28.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Protease Assay: After incubation of 28 units of α 1-3, 6 Galactosidase with 0.2 nmol of a standard mixture of proteins in a 20 μ l reaction, for 20 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

Note: Recommended storage temperature is 4°C. Avoid repeated freeze/thaw cycles

Heat Inactivation: 65°C for 10 minutes.

References:

1. McLeod, E., New England Biolabs, Inc. unpublished results.
2. Wong-Madden, S.T. and Landry, D. (1995) *Glycobiology* 5, 19–28.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.