

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	StickTogether™ DNA Ligase Buffer
Catalog Number:	B0535S
Concentration:	2 X Concentrate
Packaging Lot Number:	10114677
Expiration Date:	07/2024
Storage Temperature:	-20°C
Specification Version:	PS-B0535S v1.0
Composition (1X):	66 mM Tris-HCl, 10 mM MgCl2, 1 mM DTT, 1 mM ATP, 7.5 % PEG 6000, (pH 7.6 @ 25°C)

StickTogether™ DNA Ligase Buffer Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B0535AVIAL	StickTogether™ DNA Ligase Buffer	10113004	Pass	

Assay Name/Specification	Lot # 10114677
RNase Activity (Buffer) A 10 µl reaction in 1X StickTogether [™] DNA Ligase Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	Pass
pH (buffers/solutions) The pH of 2X StickTogether™ DNA Ligase Buffer is between pH 7.5 and 7.7 at 25°C.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Functional Testing (DNA Ligase Buffer) A 20 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 50 ng of Lambda-HindIII DNA and 2,000 units of T4 DNA Ligase incubated for 5 minutes at 25°C results in approximately >95% ligation of the DNA fragments as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking, Buffer) A 50 μl reaction in 1X StickTogether™ DNA Ligase Buffer containing 1 μg of	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10114677
supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to	
the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 12 Jul 2021

Michae m. l

Michael Tonello Packaging Quality Control Inspector 12 Jul 2021

