

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Nuclease-free Water

Catalog Number: B1500S
Packaging Lot Number: 10139011
Expiration Date: 09/2023
Storage Temperature: 25°C

Specification Version: PS-B1500S/L v2.0

| Nuclease-free Water Component List |                       |            |                      |  |
|------------------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b>             | Component Description | Lot Number | Individual QC Result |  |
| B1500SVIAL                         | Nuclease-free Water   | 10109003   | Pass                 |  |

| Assay Name/Specification  | Lot # 10139011 |
|---|----------------|
| qPCR DNA Contamination (E. coli Genomic, Water)  Nuclease-free Water is used to make a qPCR master mix and screened across a 96 well plate for the presence of E. coli genomic DNA using 40 cycles of SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Melt curve analysis results in < 5% positive samples above background. | Pass           |
| UV-Visable Scan A UV-Visible scan using a spectrophotometer that covers the range of 200nm to 800nm will have no detectable peaks above background.   | Pass           |
| RNase Activity (Extended Digestion, Water) A 10 µl reaction in 1X NEBuffer 4 containing 40 ng of RNA transcript with Nuclease-free Water is incubated at 37°C. After incubation for 16 hours, no detectable degradation of the RNA is observed as determined by gel electrophoresis using fluorescent detection.                                    | Pass           |
| Endotoxin Testing (Endosafe®) Each test channel of the cartridge is loaded with 25 μl of Nuclease-free Water, then placed into the Endosafe MCS reader for analysis resulting in a measurement of <0.01 EU/ml.  | Pass           |
| Endonuclease Activity (Nicking, Water) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 RF I DNA with Nuclease-free Water incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.  | Pass           |



B1500S / Lot: 10139011

Page 1 of 2

| Assay Name/Specification  | Lot # 10139011 |
|---|----------------|
| Non-Specific DNase Activity (16 Hour, Water)  | Pass           |
| A 50 µl reaction in CutSmart® Buffer containing 1 µg of PhiX174-HaeIII DNA with     |                |
| Nuclease-free Water incubated for 16 hours at 37°C results in a DNA pattern free of |                |
| detectable nuclease degradation as determined by agarose gel electrophoresis.       |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Michael Dalton
Production Scientist
07 Feb 2022

Corey Rabeau

Packaging Quality Control Inspector

07 Feb 2022



B1500S / Lot: 10139011

Page 2 of 2