

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: NEBuffer™ 4

Catalog Number: B7004S

Concentration: 10 X Concentrate

Packaging Lot Number: 10096784
Expiration Date: 12/2023
Storage Temperature: -20°C

Specification Version: PS-B7004S v1.0

Composition (1X): 50 mM Potassium Acetate , 20 mM Tris-acetate, 10 mM Magnesium

Acetate, 1 mM DTT, (pH 7.9 @ 25°C)

NEBuffer™ 4 Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
B7004SVIAL	NEBuffer™ 4	10089408	Pass	

Assay Name/Specification	Lot # 10096784
Conductivity (buffers/solutions)	Pass
The conductivity of 10X NEBuffer 4 is between 36 and 54 mS at 25°C.	
Endonuclease Activity (Nicking, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 4 containing 1 μg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	
determined by agarose gerelestrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 4 plus 100 μg/ml Bovine Serum Albumin containing 1	
μg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete	
digestion of the substrate DNA as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, BSA, Buffer)	Pass
A 50 µl reaction in 1X NEBuffer 4 plus 100 µg/ml Bovine Serum Albumin containing 1	1 433
µg of Lambda dam- DNA and 1 unit of Clal incubated for 1 hour at 37°C results in	
complete digestion of the substrate DNA as determined by agarose gel	
electrophoresis.	
	_
Non-Specific DNase Activity (16 hour, Buffer)	Pass
A 50 μl reaction in 1X NEBuffer 4 containing 1 μg of PhiX174-HaeIII DNA incubated	
for 16 hours at 37°C results in a DNA pattern free of detectable nuclease	



B7004S / Lot: 10096784

Page 1 of 2

Assay Name/Specification	Lot # 10096784
degradation as determined by agarose gel electrophoresis.	
pH (buffers/solutions) The pH of 10X NEBuffer 4 is between pH 7.8 and 8.0 at 25°C.	Pass
RNase Activity (Buffer)	Pass
A 10 µl reaction in 1X NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA	
remains intact as determined by fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Michael Dalton
Production Scientist

09 Feb 2021

Josh Hersey

Packaging Quality Control Inspector

09 Feb 2021



B7004S / Lot: 10096784

Page 2 of 2