

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Diluent A
Catalog Number: B8001S

Concentration: 1 X Concentrate

Lot Number: 10042753
Expiration Date: 04/2022
Storage Temperature: -20°C

Specification Version: PS-B8001S v1.0

Composition (1X): 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 µg/ml BSA

, 50 % Glycerol, (pH 7.4 @ 25°C)

| Diluent A Component List | | | | |
|--------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| B8001SVIAL | Diluent A | 10042697 | Pass | |

| Assay Name/Specification | Lot # 10042753 |
|--|----------------|
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of PhiX174-HaeIII DNA and a minimum of 10 µl of Diluent A incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 µl of Diluent A incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Diluent A is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |
| pH (buffers/solutions) The pH of 1X Diluent A is between pH 7.3 and 7.5 at 25°C. | Pass |
| qPCR DNA Contamination (E. coli Genomic) A minimum of 1 μl of Diluent A is screened for the presence of E. coli genomic DNA | Pass |



B8001S / Lot: 10042753



| Assay Name/Specification | Lot # 10042753 |
|--|----------------|
| using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results | |
| are quantified using a standard curve generated from purified E. coli genomic DNA. | |
| The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | |

This product has been tested and shown to be in compliance with all specifications.

Production Scientist

15 Apr 2019

Michael Tonello

Packaging Quality Control Inspector

18 Apr 2019