

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Diluent B
Catalog Number: B8002S

Concentration: 1 X Concentrate

Packaging Lot Number: 10098497
Expiration Date: 07/2023
Storage Temperature: -20°C

Specification Version: PS-B8002S v1.0

Composition (1X): 300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol , 500 μg/ml BSA, (pH 7.4 @ 25°C)

Diluent B Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
B8002SVIAL	Diluent B	10070597	Pass	

Assay Name/Specification	Lot # 10098497
RNase Activity (Extended Digestion) A 10 μ I reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ I of Diluent B is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 1 µl of Diluent B is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
pH (buffers/solutions) The pH of 1X Diluent B is between pH 7.3 and 7.5 at 25°C.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of PhiX174-HaeIII DNA and a minimum of 10 µl of Diluent B incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking)	Pass



B8002S / Lot: 10098497

Page 1 of 2

A FO ut reaction in CutSmort® Buffer containing 1 up of supercoiled DhiV174 DNA and	Lot # 10098497
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 µl of Diluent B incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Michael Dalton **Production Scientist**

18 May 2021

Josh Hersey

Packaging Quality Control Inspector

18 May 2021

