

New England Biolabs Certificate of Analysis

Product Name: T4 Polynucleotide Kinase
Catalog Number: M0201S
Concentration: 10,000 U/ml
Unit Definition: One Richardson unit is defined as the amount of enzyme catalyzing the incorporation of 1 nmol of acid insoluble [³³P] in 30 minutes at 37°C.
Packaging Lot Number: 10138992
Expiration Date: 12/2023
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.1 μM ATP , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0201S/L v1.0


T4 Polynucleotide Kinase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0201SVIAL	T4 Polynucleotide Kinase	10129471	Pass
B0201SVIAL	T4 Polynucleotide Kinase Reaction Buffer	10119056	Pass

Assay Name/Specification	Lot # 10138992
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of T4 Polynucleotide Kinase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) T4 Polynucleotide Kinase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of T4 Polynucleotide Kinase is screened for the presence of E.</p>	Pass

Assay Name/Specification	Lot # 10138992
<p>coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	
<p>Non-Specific DNase Activity (16 Hour) A 50 μl reaction in T4 Polynucleotide Kinase Reaction Buffer containing 1 μg of Lambda DNA and a minimum of 100 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 μl reaction in T4 Polynucleotide Kinase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 100 units of T4 Polynucleotide Kinase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 μl reaction in T4 Polynucleotide Kinase Reaction Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of T4 Polynucleotide Kinase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>DNase Activity (Labeled Oligo, 5' extension) A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>DNase Activity (Labeled Oligo, 3' extension) A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass
<p>Double Stranded DNase Activity (Labeled Oligo) A 50 μl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 50 units of T4 Polynucleotide Kinase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
15 Feb 2022



Josh Hersey
Packaging Quality Control Inspector
15 Feb 2022