

## New England Biolabs Certificate of Analysis

*Product Name:* HpaII Methyltransferase  
*Catalog #:* M0214S/L  
*Concentration:* 4,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to protect 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by HpaII restriction endonuclease.  
*Lot #:* 0181802  
*Assay Date:* 02/2018  
*Expiration Date:* 2/2019  
*Storage Temp:* -20°C  
*Storage Conditions:* 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA, 50 % Glycerol, (pH 7.4 @ 25°C)  
*Specification Version:* PS-M0214S/L v1.0  
*Effective Date:* 01 Aug 2016

Assay Name/Specification (minimum release criteria)	Lot #0181802
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Methylase Activity (dam Methylase)</b> - A 20 µl reaction in HpaII Methyltransferase Buffer supplemented with 80 µM S-adenosylmethionine containing 1 µg Lambda DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 4 hours at 37°C did not protect the DNA from digestion by MboI as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 40 units of HpaII Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> - HpaII Methyltransferase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>
<b>RNase Activity (Extended Digestion)</b> - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of HpaII Methyltransferase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



---

**New England Biolabs  
Certificate of Analysis**



---

Authorized by  
Derek Robinson  
01 Aug 2016





---

Inspected by  
Mala Samaranayake  
09 Mar 2018