

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Vent® DNA Polymerase

Catalog Number: M0254S
Concentration: 2,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10

nmol of dNTP into acid-insoluble material in 30 minutes at 75°C.

Packaging Lot Number: 10072331
Expiration Date: 04/2022
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.1 %

Triton®X-100 , 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0254S/L v1.0

Vent® DNA Polymerase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0254SVIAL	Vent® DNA Polymerase	10069192	Pass	
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10067018	Pass	
B1003SVIAL	Magnesium Sulfate (MgSO <sub>4</sub> ) Solution	10073346	Pass	

Assay Name/Specification	Lot # 10072331
Endonuclease Activity (Nicking, Polymerase, dNTP) A 50 μl reaction in ThermoPol® Reaction Buffer in the presence of 400 μM dNTPs containing 1 μg of supercoiled pUC19 DNA and a minimum of 20 units of Vent® DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
PCR Amplification (2.0 kb Lambda DNA) A 25 μl reaction in ThermoPol® Reaction Buffer in the presence of 200 μM dNTPs and 0.5 μM primers containing 5 ng Lambda DNA with 0.25 units of Vent® DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Vent® DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE)	Pass



M0254S / Lot: 10072331

Page 1 of 2

Assay Name/Specification	Lot # 10072331
Vent® DNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 2 units of Vent® DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Vent® DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Christie Vazquez Production Scientist 05 Aug 2020 Michael Tonello

Packaging Quality Control Inspector

05 Aug 2020



M0254S / Lot: 10072331