

New England Biolabs Certificate of Analysis

Product Name: RecJf
Catalog Number: M0264S
Concentration: 30,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to produce 0.05 nmol TCA soluble deoxyribonucleotide in a total reaction volume of 50 µl in 30 minutes at 37°C.
Packaging Lot Number: 10091327
Expiration Date: 12/2022
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 µg/ml BSA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0264S/L v2.0


RecJf Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0264SVIAL	RecJf	10088327	Pass
B7002SVIAL	NEBuffer™ 2	10089398	Pass

Assay Name/Specification	Lot # 10091327
<p>Ligation and Recutting (Terminal Integrity, Digested DNA) A 50 µl reaction in 1X CutSmart® Buffer containing 4 µg of pUC19-SphI digest and a minimum of 60 units of RecJf incubated for 4 hours at 37°C results in >95% ligation of the DNA fragments as determined by agarose gel electrophoresis. Of these ligated fragments, >95% can be recut with SphI.</p>	Pass
<p>Ligation and Recutting (Terminal Integrity, Digested DNA) A 50 µl reaction in 1X CutSmart® Buffer containing 4 µg of PhiX174-HaeIII digest and a minimum of 60 units of RecJf incubated for 4 hours at 37°C results in >95% ligation of the DNA fragments as determined by agarose gel electrophoresis. Of these ligated fragments, >95% can be recut with HaeIII.</p>	Pass
<p>RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 30 units of RecJf is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

Assay Name/Specification	Lot # 10091327
<p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 90 units of RecJf incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Circular Single Stranded DNA) A 50 µl reaction in CutSmart® Buffer containing 1 µg of PhiX174 Virion DNA and a minimum of 90 units of RecJf incubated for 4 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



John Greci
Production Scientist
18 Dec 2020



Michael Tonello
Packaging Quality Control Inspector
18 Dec 2020