

New England Biolabs Certificate of Analysis

Product Name: OneTaq® Hot Start 2X Master Mix with GC Buffer
Catalog Number: M0485L
Concentration: 2 X Concentrate
Packaging Lot Number: 10221835
Expiration Date: 09/2025
Storage Temperature: -20°C
Specification Version: PS-M0485S/L v2.0
Composition (1X): 80 mM Tris-SO4 (pH 9.2 @ 25°C), 20 mM (NH4)2SO4, 2 mM MgSO4, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 5 % DMSO, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 25 units/ml OneTaq® Hot Start DNA Polymerase


| OneTaq® Hot Start 2X Master Mix with GC Buffer Component List | | | |
|---|--|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M0485SVIAL | OneTaq® Hot Start 2X Master Mix with GC Buffer | 10205869 | Pass |
| B9026AVIAL | OneTaq® High GC Enhancer | 10210663 | Pass |

| Assay Name/Specification | Lot # 10221835 |
|--|----------------|
| Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs including [³ H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X OneTaq® Hot Start Master Mix with GC Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| PCR Amplification (Buffer Dependent, >65% GC-rich, Master Mix) A 25 µl reaction in 1X OneTaq® Hot Start Master Mix with GC Buffer and 0.2 µM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the buffer-dependent production of the 737 bp product. | Pass |
| PCR Amplification (Enhancer Dependent, >70% GC-rich, Master Mix) A 25 µl reaction in 1X OneTaq® Hot Start Master Mix with GC Buffer and 20% OneTaq® | Pass |

| Assay Name/Specification | Lot # 10221835 |
|---|----------------|
| <p>High GC Enhancer in the presence of 0.2 μM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the enhancer-dependent production of the 627 bp product.</p> | |
| <p>PCR Amplification (Hot Start 2 kb Lambda DNA) A 25 μl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 μM dNTPs and 0.2 μM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.</p> | Pass |
| <p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μl of OneTaq® Hot Start 2X Master Mix with GC Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.


 Lea Antonopoulos
 Production Scientist
 22 Sep 2023


 Josh Hersey
 Packaging Quality Control Inspector
 12 Feb 2024