

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: EnGen® Spy dCas9 (SNAP-tag)

Catalog Number: M0652T
Concentration: 20 µM
Unit Definition: N/A

Packaging Lot Number: 10063233
Expiration Date: 01/2022
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0652T v1.0

EnGen® Spy dCas9 (SNAP-tag) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0652TVIAL	EnGen® Spy dCas9 (SNAP-tag®)	10063232	Pass	
B7203SVIAL	NEBuffer™ 3.1	10053974	Pass	

Assay Name/Specification	Lot # 10063233
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (EnGen® Spy dCas9 (SNAP-tag®), Gel Shift Assay) A 20 µl reaction in 1X NEBuffer 3.1 containing 20 nM 100 bp FAM labeled double stranded target DNA, 20 nM TAMRA-labeled off target DNA, 100 nM sgRNA and 100 nM EnGen® Spy dCas9 (SNAP-tag®) incubated for 15 minutes at 37°C results in ≥90% binding of the substrate DNA as determined by electrophoretic mobility shift assay.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) incubated for 16 hours at 37°C results in a DNA	Pass



M0652T / Lot: 10063233

Page 1 of 2

Assay Name/Specification	Lot # 10063233
pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) EnGen® Spy dCas9 (SNAP-tag®) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 pmol of EnGen® Spy dCas9 (SNAP-tag®) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Jessica Cane Production Scientist 07 Jan 2020 Michael Tonello

Packaging Quality Control Inspector

18 Feb 2020



M0652T / Lot: 10063233

Page 2 of 2