

New England Biolabs Certificate of Analysis

Product Name: *EnGen Lba Cas12a (Cpf1)*
Catalog Number: *M0653T*
Concentration: *100 µM*
Packaging Lot Number: *10160660*
Expiration Date: *08/2024*
Storage Temperature: *-20°C*
Storage Conditions: *500 mM NaCl, 20 mM Sodium Acetate, 0.1 mM EDTA, 0.1 mM TCEP-HCl, 50% Glycerol, (pH 6.0 @ 25°C)*
Specification Version: *PS-M0653T v2.0*

EnGen Lba Cas12a (Cpf1) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0653TVIAL	EnGen® Lba Cas12a (Cpf1)	10156778	Pass
B6002SVIAL	NEBuffer™ r2.1	10156432	Pass
B0653AVIAL	EnGen® Lba Cas12a Diluent	10156821	Pass

Assay Name/Specification	Lot # 10160660
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of f-300 RNA transcript and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Functional Testing (Targeted Digestion) A 20 µl reaction in 1X NEBuffer 2.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM crRNA, and 100 nM EnGen® Lba Cas12a (Cpf1) incubated for 15 minutes at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and</p>	Pass

Assay Name/Specification	Lot # 10160660
<p>double-stranded [³H] E. coli DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p> <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda DNA and a minimum of 1 pmol of EnGen® Lba Cas12a (Cpf1) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Bhairavi Jani
Production Scientist
19 Aug 2022



Michael Tonello
Packaging Quality Control Inspector
19 Aug 2022