

New England Biolabs Certificate of Analysis

Product Name: α 1-2,3 Mannosidase
Catalog #: P0729S/L
Concentration: 32,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave > 95% of the non-reducing terminal α -D-mannose from 1 nmol Man α 1-3Man β 1-4GlcNAc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10 μ l.
Lot #: 0191711
Assay Date: 11/2017
Expiration Date: 11/2018
Storage Temp: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0729S/L v1.0
Effective Date: 19 Feb 2016

| Assay Name/Specification (minimum release criteria) | Lot #0191711 |
|---|--------------|
| Glycosidase Activity (Endo F1, F2, H) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (Endo F2, F3) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (PNGase F) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β-Mannosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Mannosidase substrate (Man β 1-4Man β 1-4Man-AMC) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β-Xylosidase) - A 10 μ l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β -Xylosidase substrate (Xyl β 1-4Xyl β 1-4Xyl β 1-4Xyl-AMC) and 32 units of α 1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |

New England Biolabs Certificate of Analysis

| Assay Name/Specification (minimum release criteria) | Lot #0191711 |
|--|--------------|
| <p>Glycosidase Activity (β1-3 Galactosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β1-4 Galactosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α-Glucosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Glucosidase substrate (Glcα1-6Glcα1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α-Neuraminidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Neuraminidase substrate (Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-2 Fucosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-2Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-3 Fucosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Fucosidase substrate (Fucα1-3Galβ1-4GlcNAcβ1-3Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |

New England Biolabs Certificate of Analysis

| Assay Name/Specification (minimum release criteria) | Lot #0191711 |
|--|--------------|
| <p>Glycosidase Activity (α1-3 Galactosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-3Galβ1-4GlcNAc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α1-6 Galactosidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Glycosidase Activity (α-N-Acetylgalactosaminidase) - A 10 μl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-N-Acetylgalactosaminidase substrate (GalNAcα1-3(Fucα1-2)Galβ1-4Glc-AMC) and 32 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p> | Pass |
| <p>Protease Activity (SDS-PAGE) - A 20 μl reaction in 1X Glyco Buffer 1 containing 24 μg of a standard mixture of proteins and a minimum of 400 units of α1-2,3 Mannosidase incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p> | Pass |
| <p>Protein Purity Assay (SDS-PAGE) - α1-2,3 Mannosidase is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> | Pass |



Authorized by
Derek Robinson
19 Feb 2016



Inspected by
Brad Landgraf
31 Oct 2017

