

## New England Biolabs Certificate of Analysis

**Product Name:**  $\beta$ 1-4 Galactosidase S  
**Catalog Number:** P0745L  
**Concentration:** 8,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to cleave > 95% of the terminal,  $\beta$ -D-galactose from 1 nmol Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc-7-amino-4-methyl-coumarin (AMC), in 1 hour at 37°C in a total reaction volume of 10  $\mu$ l.  
**Packaging Lot Number:** 10239803  
**Expiration Date:** 11/2025  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0745S/L v1.0

<b><math>\beta</math>1-4 Galactosidase S Component List</b>			
<b>NEB Part Number</b>	<b>Component Description</b>	<b>Lot Number</b>	<b>Individual QC Result</b>
P0745LVIAL	$\beta$ 1-4 Galactosidase S	10217665	<b>Pass</b>
B1727SVIAL	10X GlycoBuffer 1	10229621	<b>Pass</b>

<b>Assay Name/Specification</b>	<b>Lot # 10239803</b>
<p><b>Glycosidase Activity (Endo F1, F2, H)</b>            A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (Endo F2, F3)</b>            A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (PNGase F)</b>            A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled PNGase F substrate (Fluoresceinated fetuin triantennary) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10239803
<p><b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Glucosidase substrate (Glc<math>\alpha</math>1-6Glc<math>\alpha</math>1-4Glc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuca1-2)Gal<math>\beta</math>1-4Glc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Neuraminidase substrate (Neu5Ac<math>\alpha</math>2-3Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>1-2 Fucosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuca1-2Gal<math>\beta</math>1-4Glc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Fucosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuca1-3Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Galactosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Mannosidase)</b> A 10 <math>\mu</math>l reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-3Man<math>\beta</math>1-4GlcNAc-AMC) and 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	Pass
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Galactosidase)</b></p>	Pass

Assay Name/Specification	Lot # 10239803
<p>A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Galactosidase substrate (Galα1-6Galα1-6Glcα1-2Fru-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	
<p><b>Glycosidase Activity (α1-6 Mannosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled α-Mannosidase substrate (Manα1-6Manα1-6(Manα1-3)Man-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-Mannosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-N-Acetylgalactosaminidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylgalactosaminidase substrate (GalNAcβ1-4Galβ1-4Glc-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-N-Acetylglucosaminidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-N-Acetylglucosaminidase substrate (GlcNAcβ1-4GlcNAcβ1-4GlcNAc-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-Xylosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β1-3 Galactosidase)</b> A 10 µl reaction in Glyco Buffer 1 containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 80 units of β1-4 Galactosidase S incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Protease Activity (SDS-PAGE)</b> A 20 µl reaction in 1X Glyco Buffer 1 containing 24 µg of a standard mixture of</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10239803
<p>proteins and a minimum of 80 units of <math>\beta</math>1-4 Galactosidase S incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p> <p><b>Protein Purity Assay (SDS-PAGE)</b>  <math>\beta</math>1-4 Galactosidase S is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Alicia Bielik  
Production Scientist  
22 Nov 2023



Michael Tonello  
Packaging Quality Control Inspector  
15 Apr 2024