

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: BamHI
Catalog Number: R0136M
Concentration: 100,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10091750
Expiration Date: 10/2022
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0136T/M v1.0

BamHI Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0136MVIAL	BamHI	10086193	Pass	
B7203SVIAL	NEBuffer™ 3.1	10085493	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10084972	Pass	

Assay Name/Specification	Lot # 10091750
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of BamHI, religated and	Pass
transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of BamHl incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [3H] E. coli DNA and a minimum of 100 units of BamHl incubated for	
4 hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of Lambda DNA with BamHI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	



R0136M / Lot: 10091750

Page 1 of 2

Assay Name/Specification	Lot # 10091750
>95% can be recut with BamHI.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 20 units of BamHl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

24 Nov 2020

Josh Hersey

Packaging Quality Control Inspector

24 Nov 2020



R0136M / Lot: 10091750

Page 2 of 2