

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Xmal
Catalog Number: R0180L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

pXba in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10155954
Expiration Date: 06/2024
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0180S/L v1.0

Xmal Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0180LVIAL	Xmal	10155953	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10153339	Pass	
B6004SVIAL	rCutSmart™ Buffer	10153337	Pass	

Assay Name/Specification	Lot # 10155954
Protein Purity Assay (SDS-PAGE)	Pass
Xmal is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 50 Units of Xmal incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
donversion to the nicked form as determined by agarose ger electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 μl reaction in CutSmart [™] Buffer containing 1 μg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 50 units of Xmal incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 10 Units of Xmal incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	



R0180L / Lot: 10155954

Page 1 of 2

Assay Name/Specification	Lot # 10155954
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pXba DNA with Xmal, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95%	
can be recut with Xmal.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

15 Jul 2022

Erin Varney

Packaging Quality Control Inspector

15 Jul 2022



R0180L / Lot: 10155954

Page 2 of 2