

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Stul

Catalog Number: R0187S

Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10060532
Expiration Date: 11/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0187S/L v1.0

Stul Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0187SVIAL	Stul	10060530	Pass	
B7204SVIAL	CutSmart® Buffer	10061299	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10055732	Pass	

Assay Name/Specification	Lot # 10060532
Blue-White Screening (Terminal Integrity)	Pass
A sample of Litmus 28i vector linearized with a 10-fold excess of Stul, religated	
and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Protein Purity Assay (SDS-PAGE)	Pass
Stul is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a	
minimum of 10 Units of Stul incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
to the nicked form as determined by agarose ger electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 μl reaction in CutSmart [™] Buffer containing 1 μg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 300 units of Stul incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
nous at a constant of the total radioactivity.	



R0187S / Lot: 10060532

Page 1 of 2

Assay Name/Specification	Lot # 10060532
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Stul, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Stul.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of Stul incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis
Production Scientist

02 Dec 2019

Jay Minichiello

Packaging Quality Control Inspector

22 Jan 2020



R0187S / Lot: 10060532

Page 2 of 2