

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: PpuMI
Catalog Number: R0506S
Concentration: 10,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction

volume of 50 μl.

Lot Number: 10021254
Expiration Date: 11/2020
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0506S/L v1.0

PpuMI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0506SVIAL	PpuMI	10020324	Pass	
B7204SVIAL	CutSmart® Buffer	10021118	Pass	

Assay Name/Specification	Lot # 10021254
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 30 units of PpuMI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
ger electronical communication and against ger electrophic content	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 100 units of PpuMI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Thours at 37 O releases <0.176 or the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of pBC4 DNA with PpuMI, >95% of the DNA fragments can	
be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95%	
can be recut with PpuMI.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a	
minimum of 100 Units of PpuMI incubated for 16 hours at 37°C results in a DNA	



R0506S / Lot: 10021254

Page 1 of 2

Assay Name/Specification	Lot # 10021254
pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso Production Scientist

01 Oct 2018

Michael Tonello

Packaging Quality Control Inspector

01 Nov 2018

