

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	PfIMI
Catalog Number:	R0509S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10167243
Expiration Date:	10/2024
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R0509S/L v1.0

PfIMI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0509SVIAL	PfIMI	10167242	Pass	
B6003SVIAL	NEBuffer™ r3.1	10146827	Pass	

Assay Name/Specification	Lot # 10167243
Protein Purity Assay (SDS-PAGE) PfIMI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 μ I reaction in NEBuffer 3.1 containing 1 μ g of supercoiled pNEB193 DNA and a minimum of 50 units of PfIMI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of PfIMI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with PfIMI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with PfIMI.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10167243
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 30	
Units of PfIMI incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun

Production Scientist 12 Oct 2022

Michae 11,

Michael Tonello Packaging Quality Control Inspector 18 Oct 2022

