

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Pmll
Catalog Number: R0532S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

Lambda DNA (HindIII digest) DNA in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10098815
Expiration Date: 01/2022
Storage Temperature: -20°C

Storage Conditions: 25 mM KCl, 25 mM Tris-HCl (pH 7.5), 1 mM DTT, 0.5 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0532S/L v2.0

| PmII Component List | | | | |
|------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0532SVIAL | PmII | 10097016 | Pass | |
| B7204SVIAL | CutSmart® Buffer | 10093127 | Pass | |

| Assay Name/Specification | Lot # 10098815 |
|---|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of PmII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda HindIII DNA with PmII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PmII. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII DNA and a minimum of 100 Units of PmII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and | Pass |



R0532S / Lot: 10098815

Page 1 of 2

| Assay Name/Specification | Lot # 10098815 |
|---|----------------|
| a minimum of 20 units of PmII incubated for 4 hours at 37°C results in <20% | |
| conversion to the nicked form as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

24 Feb 2021

Michael Tonello

Packaging Quality Control Inspector

24 Feb 2021

