

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | BsaBl |
|------------------------|---|
| Catalog Number: | R0537S |
| Concentration: | 10,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (dam-) in 1 hour at 60°C in a total reaction volume of 50 μl. |
| Packaging Lot Number: | 10063042 |
| Expiration Date: | 11/2021 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 μg/ml BSA |
| Specification Version: | PS-R0537S/L v1.0 |

| BsaBI Component List | | | | |
|------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0537SVIAL | BsaBl | 10058788 | Pass | |
| B7204SVIAL | CutSmart® Buffer | 10061304 | Pass | |

| Assay Name/Specification | Lot # 10063042 |
|---|----------------|
| Non-Specific DNase Activity (16 hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda dam- DNA and a minimum of 10 units of BsaBl incubated for 16 hours at 60°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda dam- DNA with BsaBI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsaBI. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of BsaBI incubated for 4 hours at 60°C releases <0.1% of the total radioactivity. | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10063042 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE) BsaBI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist 04 Nov 2019

Michae 11.

Michael Tonello Packaging Quality Control Inspector 22 Jan 2020

