

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: BsaBI
Catalog Number: R0537S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA (dam-) in 1 hour at 60°C in a total reaction volume of

50 μl.

Packaging Lot Number: 10174704
Expiration Date: 12/2024
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 μg/ml BSA

Specification Version: PS-R0537S/L v1.0

| BsaBl Component List | | | | |
|------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0537SVIAL | BsaBl | 10174702 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10173661 | Pass | |

| Assay Name/Specification | Lot # 10174704 |
|--|----------------|
| Non-Specific DNase Activity (16 hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda dam- DNA and a minimum of 10 units of BsaBl incubated for 16 hours at 60°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme. | Pass |
| Protein Purity Assay (SDS-PAGE) BsaBI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of BsaBl incubated for 4 hours at 60°C releases <0.1% of the total radioactivity. | Pass |



R0537S / Lot: 10174704

Page 1 of 2

| Assay Name/Specification | Lot # 10174704 |
|---|----------------|
| Ligation and Recutting (Terminal Integrity) | Pass |
| After a 10-fold over-digestion of Lambda dam- DNA with BsaBI, >95% of the DNA | |
| fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated | |
| fragments, >95% can be recut with BsaBI. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń Production Scientist

04 Ján 2023

Michael Tonello

Packaging Quality Control Inspector

05 Jan 2023



R0537S / Lot: 10174704

Page 2 of 2