

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: AfIIII

Catalog Number: R0541S

Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in NEBuffer™ r3.1 in 1 hour at 37°C in a total

reaction volume of 50 μl.

Packaging Lot Number: 10225703
Expiration Date: 10/2025
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 500 mM NaCl ,1 mM DTT , 0.1 mM EDTA , 50 % Glycerol

, 200 μg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0541S/L v2.0

AfIIII Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0541SVIAL	AfIIII	10211699	Pass	
B6003SVIAL	NEBuffer™ r3.1	10221488	Pass	

Assay Name/Specification	Lot # 10225703
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of AfIIII incubated for	
4 hours at 37°C releases <0.1% of the total radioactivity.	
Functional Testing (15 minute Digest)	Pass
A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and 1 µl of AfIIII	
incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	
agaroso gor crock opriorosic.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 5-fold over-digestion of Lambda DNA with AfIIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with AfIII.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and a minimum of	
30 units of AfIIII incubated for 16 hours at 37°C results in a DNA pattern free of	



R0541S / Lot: 10225703

Page 1 of 2

Assay Name/Specification	Lot # 10225703
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) AfIIII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of AfIIII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun \

Production Scientist

24 Oct 2023

Michael Tonello

Packaging Quality Control Inspector

16 Jan 2024



R0541S / Lot: 10225703

Page 2 of 2