

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | BsmFl |
|------------------------|--|
| Catalog Number: | R0572S |
| Concentration: | 2,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μ g of pBR322 DNA in 1 hour at 65°C in a total reaction volume of 50 μ l. |
| Packaging Lot Number: | 10065391 |
| Expiration Date: | 02/2022 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA |
| Specification Version: | PS-R0572S/L v1.0 |

| BsmFI Component List | | | | |
|------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0572SVIAL | BsmFl | 10065390 | Pass | |
| B7204SVIAL | CutSmart® Buffer | 10064409 | Pass | |

| Assay Name/Specification | Lot # 10065391 |
|---|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 10 Units of BsmFl incubated for 4 hours at 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 20 units of BsmFI incubated for 4 hours at 65°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of pBR322 DNA with BsmFI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsmFI. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of pBR322 DNA and a minimum of 2 Units of BsmFl incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

This product has been tested and shown to be in compliance with all specifications.

Anthony Frankis Production Scientist 30 Jan 2020

AL

Minichiello Packaging Quality Control Inspector 11 Feb 2020

